

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Monday, 29 July

08:00 -- 10:00

Room: 206A

BM1A • Laser Processing of Optical Glasses and Poling

Presider: Martin Bernier; Université Laval, Canada and Martin Bernier; Université Laval, Canada

BM1A.1 • 08:00 Invited

Ultrafast Laser Volume Nanostructuring of Transparent Materials: From Nanophotonics to Nanomechanics, Yves Bellouard¹; ¹*Ecole Polytechnique Federale de Lausanne, Switzerland*. When exposed to ultrafast lasers, materials can be transformed in their volume, leading to intriguing nano-crystallization and sub-wavelength pattern formations. We will discuss occurrences in various glass substrates and applications in light harvesting and nano-mechanics.

BM1A.2 • 08:30

Throughput Enhancement of Type-a Volume Bragg Gratings Inscribed by Femtosecond Laser in Burst Mode for Industrial Applications, Joelle J. Harb^{1,2}, Lauris Talbot³, Yannick Petit^{2,1}, Martin Bernier³, Lionel Canioni^{2,1}; ¹*ICMCB, France*; ²*University of Bordeaux, France*; ³*University of Laval, Canada*. We present Type-A volume Bragg gratings inscribed within minutes using a 100 W laser in burst mode and the phase-mask approach. This surpasses Gaussian-Bessel methods by 38-fold while decoupling efficient thermal management from photochemical processes.

BM1A.3 • 08:45

Enhanced Light-Matter Interactions in Spherical Bragg Resonators, Yalina Garcia-Puente¹, Raman Kashyap¹; ¹*Polytechnique Montreal, Canada*. Our research optimized Spherical Bragg Resonators to engineering emissions from Er³⁺ and Eu³⁺ emitters, enhancing light-matter interaction. The study achieved significant electromagnetic field and radiative decay rate enhancements, demonstrating the potential for efficient lasing and all-dielectric nanoantenna.

BM1A.4 • 09:00

Ultrafast Laser Structuring of Scattering Optical Fibers, Léo Colliard^{2,1}, Floriane Pellerin², Geoffroy Aubry², Martiane Cabié³, Thomas Neisius³, Franck Pigeonneau⁴, Réal Vallée¹, Martin Bernier¹, Matthieu Bellec², Wilfried Blanc²; ¹*Centre Optique Photonique et Laser, Canada*; ²*Université Côte d'Azur, INPHYNI, CNRS, France*; ³*Aix Marseille Univ, CNRS, Central Marseille, FSCM, CP2M, France*; ⁴*Mines ParisTech, PSL Research University, CEMEF - Centre for Material Forming, France*. We propose here an innovative structuring process of scattering optical fibers by engineering the morphology of the nanoparticles contained inside the core with a high repetition rate femtosecond laser.

BM1A.5 • 09:15

Micrometric Patterning of Luminescence and Second Order Optical Properties of a Terbium Containing Borogermanate Magneto-Optical Glass by Thermal Poling, Juliane Resges Orives², Lia Mara Marcondes², Frederic Adamietz¹, Thierry Cardinal¹, Marcelo Nalin², Marc Dussauze¹; ¹*Université de Bordeaux / CNRS, France*; ²*UNESP, Brazil*. A thermo-electrical imprinting process on a borogermanate glass containing Tb³⁺ have permitted a micrometric

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

structuring of the oxidation state of Tb ions. Correlative luminescence, Raman and SHG microscopies demonstrate modification of the glass structure accompanying large luminescence changes and the appearance of an electro optical response.

BM1A.6 • 09:30 Invited

Poling of Silicon Nitride for 2nd Order Optical Nonlinearity, Camille-Sophie Brès¹; ¹*Ecole Polytechnique Federale de Lausanne, Switzerland*. In this talk, I will cover electric-field induced second order nonlinearities in silicon nitride waveguides and microresonators. I will show results on all-optical poling in such structures, which allows for the inscription of nonlinear gratings with periodicity controlled by the involved optical waves. I will also show results on thermally assisted electric-field poling for linear phase modulation, as an necessary advancement to bring additional nonlinear functionalities to Si photonics.

08:00 -- 10:00

Room: 205A

IM1B • IPR Opening Session - New Horizons in Integrated and Nano Photonics

Presider: Judith Su; Univ of Arizona, Coll of Opt Sciences, United States

IM1B.1 • 08:00 Invited

Optomechanical Devices Harnessing Silicon Nanostructures, Carlos A. Alonso Ramos¹; ¹*C2N-CNRS, France*. Subwavelength silicon nanostructures provide unprecedented flexibility in the control of optomechanical effects. Here, we review our recent results on the use of nanostructures for the optimization of Brillouin interactions in suspended and non-suspended optomechanical cavities.

IM1B.2 • 08:30 Invited

Programmability and AI-Assisted Design of Silicon Photonic Circuits, Joyce K. Poon¹; ¹*Max-Planck-Inst fur Mikrostrukturphysik, Germany*. Abstract not available.

IM1B.3 • 09:00 Invited

Optical Trojan Beams- Guiding Light via Lagrange Points, Mercedeh Khajavikhan¹; ¹*University of Southern California, USA*. Abstract not available.

IM1B.4 • 09:30 Invited

Diamond Photonics, Paul E. Barclay¹; ¹*University of Calgary, Canada*. Abstract not available.

08:00 -- 10:00

Room: 205C

JM1D • Radiative Cooling I (Joint SOLED/NOMA)

Presider: Alon Gorodetsky; University of California Irvine, United States

JM1D.1 • 08:00 Invited

Radiative Cooling: a new Channel From Energy Sustainability to Semiconductors, Qiaoqiang Gan¹; ¹*King Abdullah Univ of Sci & Technology, Saudi Arabia*. This talk will describe recent progresses regarding radiative cooling strategies for optoelectronics, from system design to on-chip integration. Specifically, we will discuss the considerations for passive cooling of solar panels and light emitting diodes.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

JM1D.2 • 08:30 Invited

Radiative Cooling: a Sustainable Strategy for Thermal Management and Energy

Harvesting, Lili Cai¹; ¹*University of Illinois Urbana-Champaign, USA*. This talk will present novel scalable manufacturing techniques for passive daytime radiative cooling materials, enabling diverse applications spanning from building and personal thermal management for energy savings to low-grade heat harvesting for electricity generation.

JM1D.3 • 09:00 Tutorial

Radiative Cooling: 10 Years of Daytime Radiative Cooling and the Road Ahead, Aaswath P. Raman¹; ¹*University of California Los Angeles, USA*. We introduce and review ten years of progress since the first demonstration of daytime radiative cooling. We further highlight recent optical materials-driven advances and emerging applications, concluding with a discussion of the many untapped opportunities that remain to more effectively harness the ultimate renewable thermodynamic resources: the cold of space.

08:00 -- 10:00

Room: 206B

NpM1E • Photonic Computing and Novel Phenomena

Presider: Mikko Huttunen; Tampere University, Finland

NpM1E.1 • 08:00 Invited

Towards an Alternating Photonic Ising Machine Based on Polarization Symmetry

Breaking, Stephane Coen^{1,2}; ¹*Physics, The University of Auckland, New Zealand*; ²*The Dodd-Walls Centre for Photonic and Quantum Technologies, New Zealand*. We describe progress towards the realization of a novel photonic Ising machine based on polarization symmetry breaking. A birefringent defect forces the artificial spins to alternate, roundtrip to roundtrip, conferring remarkable robustness to the system.

NpM1E.2 • 08:30

Experimental Design, Implementation, and Measurements of an Optical Ising Machine Using Polarization Symmetry Breaking

Liam Quinn^{1,2}, Yiqing Xu^{1,2}, Julien Fatome³, Stuart Murdoch^{1,2}, Miro Erkintalo^{1,2}, Stephane Coen^{1,2}; ¹*University of Auckland, New Zealand*; ²*The Dodd-Walls Centre for Photonic and Quantum Technologies, New Zealand*; ³*Université Bourgogne-Franche-Comté, France*. We experimentally demonstrate a novel optical Ising machine utilizing spontaneous polarization symmetry breaking in a driven Kerr resonator constructed from off-the-shelf telecommunications optical fiber. Spin states are encoded onto two robust, easily measurable polarization states.

NpM1E.3 • 08:45

Phase Resetting in the Yamada Model of a Q-Switching Laser, Jacob Ngaha^{1,2}, Neil Broderick^{1,2}, Bernd Krauskopf^{1,2}; ¹*The University of Auckland, New Zealand*; ²*The Dodd Walls Centre for Quantum and Photonic Technologies, New Zealand*. We investigate the phase resetting of a periodic orbit of a self-pulsing laser, described by the Yamada model. We show how the return to the periodic orbit is affected by a brief perturbation to the associated oscillation.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

NpM1E.4 • 09:00 Invited

Photonic Computation Enabled by Sound Waves, Birgit Stiller¹; ¹*Max-Planck-Institute, Science of Light, Germany*. We experimentally demonstrate building blocks of photonic neural network based on volatile traveling acoustic waves. We implement an optoacoustic recurrent operator and a nonlinear activation function based on stimulated Brillouin scattering.

NpM1E.5 • 09:30

Mobility-Induced Light Localization in an Adverse Nonlinear Environment, Siyu Li¹, Juan Wu¹, Yuhui Zhuang¹, Yi Hu¹, Jingjun Xu¹; ¹*Nankai University, China*. We report a mobility-induced localization of a beam that tends to delocalize at rest in a fluid of light. Such a transition occurs as the beam catches up with intensity holes emerged in the fluid.

NpM1E.6 • 09:45

Withdrawn

08:00 -- 10:00

Room: 2101

NeM1C • Quantum Networks and Secure Networks

Presider: Marco Ruffini; Trinity College Dublin

NeM1C.1 • 08:00 Invited

The Art of Possible With Quantum Networking, Reza Nejabati¹; ¹*University of Bristol, United Kingdom*. Abstract not available.

NeM1C.2 • 08:30 Invited

Telecom Compatibility of Quantum Key Distribution, Rui Lin¹; ¹*Chalmers Tekniska Högskola, Sweden*. Abstract not available.

NeM1C.3 • 09:00 Invited

High-Rate CV-QKD Systems Leveraged by Advanced Coherent Detection Technology, Amirhossein Ghazisaeidi¹; ¹*Nokia Bell Labs France, France*. We review basic concepts of continuous-variable quantum key distribution, and report on some recent demonstrations of high secret key rate systems based on advanced modulation formats and coherent detection.

NeM1C.4 • 09:30 Invited

Planning Strategy Towards Secure Optical Core Networks, Carmen Mas Machuca¹; ¹*Universität der Bundeswehr München, Germany*. Network operators are considering the increase of the data transmission security by the use of QKD devices. We present best planning strategies to reduce the required investments while maximizing the key capacity.

08:00 -- 10:00

Room: 2104

SpM1G • Advanced Signal Processing

Presider: Georg Rademacher; Universität Stuttgart, Germany

SpM1G.1 • 08:00 Invited

Digital Signal Processing for Very-High Speed IM/DD Optical Communications, Vivian Xi Chen¹; ¹*Nokia Bell Labs, USA*. Abstract not available.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

SpM1G.2 • 08:30

Sub-Rate Sampled, Non-Integer Fractionally Spaced Volterra Nonlinear Equalizer for IM/DD Systems, Jaeyoon Kim¹, Hoon Kim¹; ¹*KAIST, Korea (the Republic of)*. We propose and demonstrate a sub-rate sampled (<2 sample/symbol), non-integer fractionally spaced Volterra equalizer for intensity-modulation/direct-detection systems. This equalizer does not require digital upsampling at the receiver DSP, and thus greatly relieves the complexity.

SpM1G.3 • 09:00 Tutorial

Title to be Announced, Sebastian Randel¹; ¹*Karlsruher Institut für Technologie, Germany*. Abstract not available.

08:00 -- 10:00

Room: 207

SoM1F • Novel Fiber Materials and Tapered Fibers

Presider: Alexander Heidt, Switzerland

SoM1F.1 • 08:00 Invited

Ice Microfibers for Optical Waveguiding, Xin Guo¹, Peizhen Xu¹, Bowen Cui¹, Xiangzheng Li¹, Limin Tong¹; ¹*Zhejiang University, China*. We succeeded in fabricating small-molecule ice microfibers with smooth surfaces and uniform diameters, and demonstrated that the microfibers can be operated for low-loss optical waveguiding and nanophotonic applications.

SoM1F.2 • 08:30 Invited

Design Strategies and Applications of Reshapable 4D Polymer Optical Fibers, Clement Strutynski¹, Frédéric Désévéday¹, Grégory Gadret¹, Claire-Hélène Brachais¹, Bertrand Kibler¹, Frédéric Smektala¹; ¹*Universite de Bourgogne, France*. The subject of the present work is the elaboration and application of shape-memory optical fibers from the thermal stretching of additively manufactured preforms.

SoM1F.3 • 09:00

Fabrication of Nanodiamond-Doped Silica Fiber With Highly Dispersive Nonlinearity, Pascal Hänzi¹, Grzegorz Stepniewski², Adam Filipkowski², Sara Lukasik³, Tomasz Kardas⁴, Yuriy Stepanenko⁵, Maciej Glowacki⁶, Mariusz Mrozek⁷, Adam Wojciechowski⁷, Valerio Romano¹, Robert Bogdanowicz⁶, Katarzyna Krupa⁵, Ryszard Buczynski³, Alexander M. Heidt¹, Mariusz Klimczak³; ¹*Institute of Applied Physics, University of Bern, Switzerland*; ²*Lukasiewicz Research Network, Poland*; ³*Faculty of Physics, University of Warsaw, Poland*; ⁴*Fluence Sp. z o.o, Poland*; ⁵*Institute of Physical Chemistry, Polish Academy of Sciences, Poland*; ⁶*Gdansk University of Technology, Poland*; ⁷*Jagiellonian University in Kraków, Poland*. We fabricate a nanodiamond-doped silica step-index fiber exhibiting a highly dispersive nonlinearity across near-infrared wavelengths. This is achieved without altering chromatic dispersion, which resembles SMF-28 fiber.

SoM1F.4 • 09:30

10-W-Level CW Waveguiding in a Subwavelength-Diameter Silica Microfiber, Jianbin Zhang¹, Xin Guo¹, Limin Tong¹; ¹*Zhejiang University, China*. Here we report low-loss continuous-wave optical waveguiding in a subwavelength-diameter silica microfiber with power

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

up to 13 W, making it favorable for high-speed optomechanical driving of microparticles and high-efficiency second/third harmonic generation.

SoM1F.5 • 09:45

Tapered Fiber With Dual Concentric Cores for Broadband Dispersion Compensation, Wenpu Geng¹, Zhi Zeng², Lin Zhang⁴, Zhongqi Pan³, Yang Yue²; ¹*Nankai University, China*; ²*Xi'an Jiaotong University, China*; ³*University of Louisiana at Lafayette, USA*; ⁴*Tianjin University, China*. A tapered fiber with two Ge-doped concentric cores is proposed to achieve flexible and slope-controllable broadband flat negative dispersion. The dispersion curve of the fundamental mode features $<0.54 \text{ ps}/(\text{nm}\cdot\text{km})$ variation from 1440 to 1700 nm.

10:30 -- 12:30

Room: 2000A

JM2A • Introductory Remarks and Plenary Session I

JM2A.1 • 10:30 Plenary Submission

Thermal Photonics and Its Implications, Shanhui Fan¹; ¹*Stanford University, USA*. We review the use of photonic structures to control thermal radiation and the implications of such control in renewable energy.

JM2A.2 • 10:30 Plenary Submission

Petascale Photonic Connectivity for Energy-Efficient Computing, Keren Bergman¹; ¹*Columbia University, USA*. Abstract not available.

14:00 -- 16:00

Room: 206A

BM3A • Laser Direct Writing in Optical Materials

Presider: Michael Withford; Macquarie University, Australia

BM3A.1 • 14:00 Invited

Internal Laser Structuring of Semiconductors: Opening a Route to Three-Dimensional Silicon Nano-Photonics, Onur Tokel¹; ¹*Bilkent University, Turkey*. Nano-fabrication in silicon is limited to surface. We use structured lasers and preformed seeds to establish controlled volumetric nano-fabrication; with feature sizes down to 100 nm, leading to nano-photonics applications in Si.

BM3A.2 • 14:30

Investigation of Ultrashort Pulse Written Higher Order VBG With Extended Apertures, Malte P. Siems², Daniel Richter², Ria G. Krämer², Georg R. Schwartz², Stefan Nolte^{2,1}; ¹*Fraunhofer Institute for Applied Optics and Precision Engineering IOF, Germany*; ²*Institute of Applied Physics, Friedrich-Schiller-Univ., Germany*. In this paper we investigate higher order femtosecond written VBG in fused silica. Focus of this investigation will be on the influence of the Bragg order on the properties across the aperture of the VBG.

BM3A.3 • 14:45

High Order Fiber Bragg Gratings Using Point-by-Point Femtosecond Laser Technique, Romain Cotillard¹, Nicolas Roussel¹; ¹*CEA, France*. Here we describe the inscription of high-order Bragg gratings. These optical components feature the combined properties of Bragg

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

gratings and Fabry-Perot cavities, open new possibilities both for sensor applications and as optical components, such as wavelength references or fiber-optic verniers.

BM3A.4 • 15:00

Femtosecond Laser-Induced Bragg Grating Waveguide Through Selective Control of Pulse Numbers, Foroogh Jafari¹, Jean-Sébastien Boisvert¹, Raman Kashyap^{1,2}, Sébastien Loranger¹; ¹*Electrical engineering, Polytechnique Montreal, Canada*; ²*Physics engineering, Polytechnique Montreal, Canada*. Single-step writing of Bragg grating waveguide is obtained by externally modulating the fs laser pulse picker using a waveform generator, which delivers a controlled number of pulses to induce refractive-index changes at each sub-Bragg period.

BM3A.5 • 15:15

Writing of Fiber Bragg Gratings With Focused fs Pulses Using a Two-Phase Mask Interferometer, Francois Ouellette¹; ¹*Chengdu University, China*. A 2-mask interferometer is used to write fiber Bragg gratings with index modulations up to 1.6×10^{-3} without H₂ loading on time scales of 1 minute, using 343 nm, 7 μJ energy, 237 fs pulses at 60 kHz repetition rates.

BM3A.6 • 15:30 Invited

Fabrication of Thin Film Lithium Niobate Electro-Optic Devices Using a Femtosecond Laser, Ya Cheng^{1,2}; ¹*Shanghai Institute of Optics and Fine Mechanics, China*; ²*East China Normal University, China*. We report recent progresses in thin film lithium niobate integrated photonics technology, in which high-performance active and passive photonic devices are uniquely fabricated using femtosecond laser direct writing.

14:00 -- 16:00

Room: 206B

JM3E • Quantum Photonics (Joint IPR/NP)

Presider: To be determined

JM3E.1 • 14:00 Invited

Frequency-Domain Quantum and Nonlinear Nanophotonics in Low-Loss Silicon Nitride, Avik Dutt¹; ¹*University of Maryland at College Park, USA*. Abstract not available.

JM3E.2 • 14:30

Time-bin Entangled Photons for Scalable Quantum Information Processing, Stefania Sciara¹, Hao Yu^{1,2}, Mario Chemnitz^{1,4}, Monika Monika³, Farzam Nosrati¹, Agnes George¹, Nicola Montaut¹, Bennet Fischer^{1,4}, Benjamin Crockett¹, Robin Helsten¹, Benjamin Wetzel⁵, Thorsten A. Goebel⁶, Ria G. Krämer⁷, Brent Little⁸, Sai Chu¹¹, Stefan Nolte^{7,6}, Zhiming Wang², José Azaña¹, William J. Munro⁹, David J. Moss¹⁰, Ulf Peschel³, Rosario Lo Franco¹², Roberto Morandotti¹; ¹*Institut national de la recherche scientifique -Centre Énergie, Matériaux et Télécommunications (INRS-EMT), Canada*; ²*Shimmer Center, Tianfu Jiangxi Laboratory, China*; ³*Institute of Solid State Theory and Optics, Friedrich Schiller University, Germany*; ⁴*Leibniz Institute of Photonic Technology, Germany*; ⁵*XLIM Research Institute, Université de Limoges, France*; ⁶*Fraunhofer Institute for Applied Optics and Precision Engineering IOF, Center of Excellence in Photonics, Germany*; ⁷*Friedrich-Schiller-University, Abbe Center of Photonics, Institute of Applied Physics, Germany*; ⁸*QXP Technology Inc., China*; ⁹*Okinawa Institute of Science and Technology Graduate University, Japan*; ¹⁰*Optical Sciences Centre, Swinburne University of Technology, Australia*; ¹¹*Department of Physics, City University of Hong Kong, Hong Kong*; ¹²*Universita di*

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Palermo, Italy. Encoding information in photonic time bin enables quantum technologies compatible with both integrated and fiber frameworks. Here, we demonstrate time-bin entangled qudits in a programmable photonic chip and in a fully fibered coupled loop system.

JM3E.3 • 14:45

Micro-Ring Resonator in Si₃N₄ as a Photon Source for Quantum Repeaters, Juan S. Durán-Gómez^{1,2}, Roberto Ramírez Alarcón¹, Mauricio Gomez Robles², Patricia Marisol del Carmen Tavares Ramirez^{1,2}, Gerardo de Jesús Rodríguez Becerra^{1,2}, Erasto Ortíz Ricardo³, Rafael Salas Montiel²; ¹*Quantum Photonics, Centro de Investigaciones en Óptica A.C., Mexico*; ²*Laboratory Light, nanomaterials, and nanotechnologies, Université de Technologie de Troyes, France*; ³*División de Ciencias e Ingenierías campus León, Universidad de Guanajuato, Mexico*. We report an integrated photon source designed in a micro-ring resonator in Si₃N₄, that produces photons capable of interacting with a quantum memory (Y₂SiO₅:Pr³⁺-crystal) by analyzing its spectral properties through the Joint Spectral Intensity.

JM3E.4 • 15:00

A Brillouin-Pumped, Four-Wave Mixing, Photon Pair Source, Alex I. Flint¹, Rex Bannerman¹, James Gates¹, Paolo Menna¹, Peter G. Smith¹; ¹*Optoelectronics Research Centre, University of Southampton, United Kingdom*. A novel method of generating photon pairs via spontaneous four-wave mixing in SMF-28 is demonstrated. We use a CW-pumped Brillouin laser where the intracavity field provides the pump power. Source characterisation will be presented.

JM3E.5 • 15:15

Quantum Squeezing in on-Chip Microring Measured With External Local Oscillator, Andrei Danilin^{1,2}, Dmitry A. Chermoshentsev^{1,3}, Anatoly V. Masalov^{1,4}, Igor A. Bilenko^{1,2}; ¹*Russian Quantum Center, Russian Federation*; ²*Faculty of Physics, Lomonosov Moscow State University, Russian Federation*; ³*Skolkovo Institute of Science and Technology, Russian Federation*; ⁴*Lebedev Physical Institute, Russian Academy of Sciences, Russian Federation*. We report the observation of 2 dB squeezed vacuum in an on-chip degenerate optical parametric oscillator in Si₃N₄ microring resonator. We experimentally show the possibility of squeezed states measurement in balanced heterodyne scheme.

14:00 -- 16:00

Room: 205C

JM3D • Radiative Cooling II (Joint SOLED/NOMA)

Presider: Aaswath Raman; University of California Los Angeles, United States

JM3D.1 • 14:00 Invited

Vapor Condensation With Daytime Radiative Cooling, Zongfu Yu¹; ¹*University of Wisconsin - Madison, USA*. Abstract not available.

JM3D.2 • 14:30

Electrochemically Active Metasurfaces for Multispectral Radiative Heat Management, Po-Chun Hsu¹; ¹*Pritzker School of Molecular Engineering, University of Chicago, USA*. The combination of reversible electrochemical reaction and metasurface allows multifunctional and multispectral light and heat management. This talk will introduce two examples, conjugated

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

polymers and metal, and the design principle to accomplish various dynamic thermoregulation applications.

JM3D.3 • 15:00

Adaptive Radiative Coolers May Not Be a Sustainable Option for Thermoregulating Buildings, Jyotirmoy Mandal¹, Jyothis Anand², Nithin J. Varghese¹; ¹*Princeton University, USA*; ²*Oak Ridge National Laboratory, USA*. Adaptive radiative coolers are more energy-efficient than traditional radiative coolers because they do not overcool buildings in the winter. However, because this entails trapping heat on earth, they may not be sustainable for building thermoregulation.

JM3D.4 • 15:15

Squid-Inspired Materials With Tunable Heat-Managing Properties, Aleksandra Strzelecka², Sanghoon Lee¹, Alon Gorodetsky¹, Panyiming Liu¹; ¹*University of California Irvine, USA*; ²*Chemical and Biomolecular Engineering, university of California, Irvine, USA*. Heat management is critical for the operation of many modern technologies. We have developed squid-skin-inspired thermoregulatory composite materials and further endow them with breathability, washability, and fabric compatibility. Our materials appear suitable for wearable applications.

JM3D.5 • 15:30 Invited

Hierarchical-Morphology Metafabric for Passive Thermal Management, Guangming Tao¹; ¹*Huazhong Univ of Science and Technology, China*. Abstract not available.

14:00 -- 16:00

Room: 205A

NpM3B • Ultrafast and Nonlinear Phenomena

Presider: Sylvie Lebrun; Centre National Recherche Scientifique, France

NpM3B.1 • 14:00 Invited

Femtosecond Fieldoscopy: Advancing Label-Free Spectroscopy and Imaging, Hanieh Fattahi¹; ¹*Max-Planck-Inst Physik des Lichts, Germany*. This presentation introduces femtosecond fieldoscopy, a cutting-edge technique that achieves super-resolution spectro-microscopy with unprecedented sensitivity and attosecond temporal resolution. It holds promise for label-free exploration of ultrafast phenomena at the nanoscale.

NpM3B.2 • 14:30

Spontaneous Symmetry Breaking in a Coupled Photonic Crystal Dimer With Periodic Driving, Rodrigues Bitha¹, Kevin Stitely¹, Bernd Krauskopf¹, Neil Broderick¹; ¹*University of Auckland, New Zealand*. This study examines the interactions between two optical modes in a periodically driven photonic crystal dimer. We find that the system features spontaneous symmetry breaking of different types of solutions.

NpM3B.3 • 14:45

Exploiting Time Reversal Symmetry for Beam Shaping in Multimode Nonlinear Fiber Optics, Arnaud Poisson¹, Alessandro Tonello¹, Christine Restoin¹, Vincent Couderc¹; ¹*XLIM Research Institute, France*. We numerically study the nonlinear propagation in multimode GRIN

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

fibers. By exploiting the time-reversal symmetry, we identify the conditions for robust on-demand modal distributions. We extend our study to the reversal of spatial beam self-cleaning.

NpM3B.4 • 15:00

Four Coherent Dispersive Waves Emission for OAM_{3,1} Mode in a Ring-Core Fiber, Wenpu Geng¹, Yuxi Fang¹, Zhi Zeng², Changjing Bao³, Zhongqi Pan⁴, Yang Yue²; ¹*Nankai University, China*; ²*Xi'an Jiaotong University, China*; ³*University of Southern California, USA*; ⁴*University of Louisiana at Lafayette, USA*. A ring-core fiber is designed to generate four DWs for OAM_{3,1} mode. Pumping under normal dispersion, the simulated coherent output spectrum spans 1100 nm with four distinct peaks at 760, 1130, 1430 and 1685 nm.

NpM3B.5 • 15:15

Exploiting the Up-Conversion Luminescence of Material Defects to Characterize Graded-Index Optical Fibers: From Silica to Soft-Glasses, Mario Ferraro^{2,3}, Fabio Mangini¹, Raffaele Filosa², Pedro Parra-Rivas¹, Yifan Sun¹, Wasyhun A. Gemechu¹, Alessandro Falaschi¹, Grzegorz Stepniewski⁴, Adam Filipkowski^{4,5}, Ryszard Buczynski^{4,5}, Vincent Couderc⁶, Stefan Wabnitz¹; ¹*DIET, University of Rome "La Sapienza", Italy*; ²*Department of Physics, University of Calabria, Italy*; ³*Nanotec, CNR, Italy*; ⁴*Faculty of Physics, University of Warsaw, Poland*; ⁵*Institute of Microelectronics and Photonics, Lukasiwicz Research Network, Poland*; ⁶*XLIM, UMR CNRS 7252, University of Limoges, France*. Drawing optical fibers intrinsically produces material defects, which are detrimental to most applications. Here we show that material defects may also be instrumental for characterizing the refractive index profile of graded-index fibers via up-conversion luminescence.

NpM3B.6 • 15:30

Multifrequency Nonlinear Pulse Propagation, David Castello-Lurbe¹, Enrique Silvestre¹, Miguel V. Andrés¹; ¹*Universitat de Valencia, Spain*. The nonlinear coefficient dependence on multiple frequencies is rigorously incorporated into the propagation equation so that the resulting nonlinear term is still straightforwardly computed. Readily observable consequences due to this multifrequency dispersion are predicted.

NpM3B.7 • 15:45

Efficient Emission From a Spintronic THz Emitter Based on Pump Distribution and Exposure, Gabriel Gandubert¹, Joel E. Nkeck¹, Xavier Ropagnol^{1,2}, Denis Morris³, Francois Blanchard¹; ¹*École de technologie supérieure (ÉTS), Canada*; ²*INRS-EMT, Canada*; ³*Physics, Université de Sherbrooke, Canada*. Efficiency of Spintronic terahertz emitters are thermally influenced by laser pulses. Using an oscillator laser, we show that adjusting the laser pump's spatial distribution and exposure time significantly increases its generation efficiency.

14:00 -- 16:00

Room: 2101

NeM3C • Short Reach and Data-Center Networks

Presider: Lena Wosinska; Chalmers Tekniska Högskola, Sweden

NeM3C.1 • 14:00 Invited

Novel Coherent Detection Schemes for Short Reach and Datacenter Networks, Budsara Boriboon¹, Ruben S. Luis¹, Benjamin J. Puttnam¹, Satoshi Shinada¹, Hideaki Furukawa¹; ¹*NICT, Japan, Japan*. This talk discusses the self-homodyne coherent detection for short-reach

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

applications. This approach can utilize a low-cost DFB laser and provides the feasibility of using an incoherent light source with multi-level modulation formats.

NeM3C.2 • 14:30 Invited

High Symbol Rate Short-Reach Communications With Silicon Photonics Components, Oskars Ozolins¹; ¹*Riga Technical University, Latvia*. Abstract not available.

NeM3C.3 • 15:00 Invited

Analog Fronthaul and Coherent Joint Transmission Technologies and Demonstrations for Next-Generation Cell-Free MIMO RAN Scenarios, Xiaodan Pang¹; ¹*KTH Royal Inst. of Technology, Sweden*. Abstract not available.

NeM3C.4 • 15:30

Optical-Amplification-Free 245/140 Gbaud OOK/PAM4 C-Band SiP Ring Resonator Modulator-Based Links, Armands Ostrovskis¹, Toms Salgals¹, Michael Koenigsmann², Kristaps Rubuls¹, Azra Farid², Benjamin Krüger², Arvids Sedulis¹, Fabio Pittala², Ryan P. Scott³, Hansjoerg Haisch², Lu Zhang⁴, Xianbin Yu⁴, Rafael Puerta⁵, Sandis Spolitis¹, Richard Schatz⁶, Katia Gallo⁶, Markus Gruen², Hadrien Louchet², Robert Jahn², Kazuo Yamaguchi², Vjaceslavs Bobrovs¹, Xiaodan Pang^{6,7}, Oskars Ozolins^{1,7}; ¹*Riga Technical University, Latvia*; ²*Keysight Technologies Deutschland GmbH, Germany*; ³*Keysight Technologies, Inc., USA*; ⁴*Zhejiang University and Zhejiang Lab, China*; ⁵*Ericsson Research, Sweden*; ⁶*KTH Royal Institute of Technology, Sweden*; ⁷*RISE Research Institutes of Sweden AB, Sweden*. We demonstrate an optical-amplification-free 245 Gbaud OOK and 140 Gbaud PAM4 links using a C-band SiP RRM-based transmitter with performance below the 6.25% overhead hard-decision forward error correction threshold after 100 meters of SMF.

14:00 -- 16:00

Room: 2104

SpM3G • Coherent Technologies I

Presider: Werner Rosenkranz; Kiel University, Germany

SpM3G.1 • 14:00 Invited

Scenarios for DSP Evolution in Long Haul Optical Communication Systems, Domaniç Lavery¹, Siddharth Varughese², Pierre Mertz², Han Sun¹; ¹*Infinera Canada Inc., United Kingdom*; ²*Infinera Corporation, USA*. We discuss how changing market requirements and the availability of coherent pluggable modules present challenges for DSP development. Possible development paths for DSP in coherent, long-haul transmission systems are discussed.

SpM3G.2 • 14:30

Fully Generalized Machine Learning-Based Equalization in Coherent Optical Transmission, Samuel Lennard¹, Fabio A. Barbosa¹, Filipe Ferreira¹; ¹*University College London, United Kingdom*. We introduce a novel training paradigm for machine learning-based equalization without any online training for dual-polarization IQ-modulated signals. Lab transmission of 30Gbaud DP-16-QAM has shown this equalizer matching conventional DSP over a range of conditions.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

SpM3G.3 • 14:45 Invited

Unlocking the Potential of Ultrawide O-Band Coherent DWDM Fiber Transmission, Yuta Wakayama¹, Daniel J. Elson¹, Filippos Balasis¹, Shohei Beppu¹, Noboru Yoshikane¹, Takehiro Tsuritani¹; ¹*KDDI Research, Japan*. This presentation delves into our latest research and hurdles in creating high-capacity O-band transmission systems, facing greater nonlinear interference and fiber attenuation than the C-band, with non-consistent behavior across wavelengths.

SpM3G.4 • 15:15

On Link Budget of PCS-Based Coherent Transceivers With Different Client Framing in C&L-Band Networks, Ahmad Abdo¹, Willy Georges², Shahab Oveis Gharan¹, Ahmed A. Omar²; ¹*Ciena Canada Inc., Canada*; ²*Zain OmanTel International, Oman*. Improvement in performance of Ethernet (ETH) versus Optical Transport Network (OTN), in probabilistic shaped constellation (PCS)-based coherent transceivers, are presented. As well, we simulated the impact of fiber characteristics on the end-of-life (EoL) capacity of a C&L-Band open line system (OLS) and Ciena's WaveLogic5 Extreme (WL5E).

SpM3G.5 • 15:30 Invited

Unlocking the Shaping Gain in Unamplified Coherent Links, Beatriz Oliveira^{5,2}, Manuel Neves^{5,2}, Jorge Silva⁵, Fernando P. Guiomar¹, Maria d. Medeiros^{4,3}, Paulo M. Monteiro^{2,5}; ¹*Instituto De Telecomunicacoes, Portugal*; ²*Universidade de Aveiro, Portugal*; ³*Instituto de Telecomunicações, Portugal*; ⁴*Universidade de Coimbra, Portugal*; ⁵*Instituto de Telecomunicações, Portugal*. The benefits of probabilistic constellation shaping in unamplified links are still under debate, putting into question Maxwell-Boltzmann (MB)-based PCS. We optimize the probability mass function and demonstrate gains of 4.2 dB at 400 Gbit/s.

14:00 -- 16:00

Room: 207

SoM3F • Hollow-core Fibers

Presider: Benoît Beaudou; GLO Photonics, France

SoM4F.1 • 14:00

Multi-Mode Deep Ultraviolet Hollow Core Fibre, Kerriane Harrington¹, Robbie Mears¹, James M. Stone¹, William J. Wadsworth¹, Jonathan C. Knight¹, Tim A. Birks¹; ¹*University of Bath, United Kingdom*. We report a multi-mode hollow core fibre, with a 33 μm diameter core, that guides ultraviolet light (330-440 nm in UV-A and 220-255 nm in UV-C). One cane stage is used for high yield fabrication.

SoM4F.2 • 14:30

Investigation Into the use of gas Permeation to Control the gas Composition and Pressure Within Hollow-Core Fibers, Kavitha Srinivasan¹, Thomas Kelly¹, Somarpita Pradhan¹, Ian Davidson¹, Radan Slavik¹, Peter Horak¹, Natalie V. Wheeler¹; ¹*University of Southampton, United Kingdom*. We investigate, both via simulations and experiments, a new approach for gas purging in hollow-core fibers based on side permeation of helium gas into the fiber's microstructure and subsequent pressure-driven flow along the fiber core

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

SoM4F.3 • 15:00

Mode Coupling and Ultimate Loss Limit in Hollow Core Fibers, Federico Melli¹, Kostiantyn Vasko², Lorenzo Rosa¹, Fetah Benabid^{2,3}, Luca Vincetti¹; ¹*Univ degli Studi Modena e Reggio Emilia, Italy*; ²*GLOPHOTONICS, France*; ³*XLIM, France*. A theoretical model describing the modes coupling in hollow core inhibited coupling fibers is presented. This model gives new insights about the ultimate limits in terms of loss and bandwidth of this kind of fibers.

SoM4F.4 • 15:15

Bending and Temperature Dependence of Polarization Mode Dispersion in Nodeless Antiresonant Hollow Core Fibers, Austin Taranta¹, Seyed Mohammad Abokhamis Mousavi¹, Eric Numkam Fokoua², Gianluca Guerra¹, Gregory Jasion¹, Konstantin Vidiajev¹, Hesham Sakr², John Hayes¹, Thomas Bradley³, Ghafour A. Mahdiraji¹, Jaroslaw Rzegocki¹, Ian Davidson¹, Radan Slavik¹, Francesco Poletti^{1,2}; ¹*Optoelectronics Research Centre, University of Southampton, United Kingdom*; ²*Microsoft Azure Fiber, United Kingdom*; ³*High-Capacity Optical Transmission Laboratory, Eindhoven University of Technology, Netherlands*. We identify empirical thermal and bending attributes of PMD in short nodeless antiresonant fibers (ARFs). Surprisingly, bend-scaling properties of ARF are comparable to solid fiber, albeit with greater magnitude, while temperature dependence is more varied.

SoM4F.5 • 15:30

Sealing Purged Mid-Infrared Hollow-Core Fibers, Qiang Fu¹, Thomas Kelly¹, Jing Meng¹, Yongmin Jung¹, Francesco Poletti¹, Natalie V. Wheeler¹, Ian Davidson¹; ¹*University of Southampton, United Kingdom*. We report two practical methods for sealing purged mid-infrared (3-4.6 μm) anti-resonant, hollow-core fibers and demonstrate that inert gas purging significantly reduces loss, which is effectively maintained by sealing.

SoM4F.6 • 15:45

Transverse Roughness: Modeling and Effects Analysis on Inhibited Coupling Fibers, Federico Melli¹, Kostiantyn Vasko², Lorenzo Rosa¹, Fetah Benabid^{3,2}, Luca Vincetti¹; ¹*Univ degli Studi Modena e Reggio Emilia, Italy*; ²*glophotonics, France*; ³*xlim, France*. A Transverse Roughness theoretical model based on the Azimuthal Fourier Decomposition is proposed to analyse the effects of this perturbation on the Confinement Loss of Hollow-Core Inhibited Coupling Fibers. Scaling laws are also given

16:30 -- 18:30

Room: 206A

BM4A • Symposium on Optical Fiber Sensors for Extreme Environments I

Presider: Guillaume Laffont

BM4A.1 • 16:30 Invited

Fiber Sensors in Tokamak for Nuclear Fusion, Jonathan Gaspar¹; ¹*Aix-Marseille Université, France*. Abstract not available.

BM4A.2 • 17:00

Survival of Nanogratings in High Melting Oxide Glasses Upon Thermal Annealing, Imane Ktafi¹, Qiong Xie¹, Gözden Torun², Yves Bellouard², Maureen Yembele³, Mathieu Allix³, Thomas Wade Hawkins⁴, John Ballato⁴, Maxime Cavillon¹, Matthieu Lancry¹; ¹*Universite Paris-Saclay, France*; ²*EPFL, Switzerland*; ³*CEMHTI – CNRS, France*; ⁴*Clemson University, USA*.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Nanogratings were fabricated using a fs-laser inside more than 20 oxide glasses. Glass viscosity mostly dictates their thermal stability but deviation from this trend was observed in glasses containing large amount of Al_2O_3 or ZrO_2

BM4A.3 • 17:15

Ceramic-Coated Type III Femtosecond Fiber Bragg Grating for High Temperature

Environments, Laure Lago¹, Guillaume Laffont¹, Rémy Bernard²; ¹CEA, France; ²Laboratoire PhLAM, Université de Lille, France. Boron nitrite-based ceramic coating is developed to protect fiber Bragg gratings (FBG) for harsh environment use. Coated FBGs are manufactured.

Characterization and tests are made in Air at different high temperature levels (700°C, 800°C, 900°C) during 500 hours. Long-term behavior (up to 1500 hours) is also studied at 800°C.

BM4A.4 • 17:30

Ultra-High Temperature Sensor Utilizing an Intrinsic Sapphire Fabry-Pérot Interferometer,

Alexander Roehrl¹, Andrea Stadler¹, Fabian Buchfellner¹, Simon Zehetmair¹, Johannes Roths¹; ¹Munich University of Applied Sciences, Germany. We propose and present a high temperature fiber optic sensor based on an intrinsic sapphire Fabry-Pérot interferometer and a single mode interrogation system for temperature measurements up to 1200°C.

BM4A.5 • 17:45 Invited

Recent Developments in Fiber Optic Sensing for Energy Infrastructure Applications,

Paul Ohodnicki¹, Khurram Naeem¹, Pengdi Zhang¹, Yang-Duan Su¹, Dolendra Karki¹, Nageswara Lalam^{2,3}, Ruishu Wright³; ¹University of Pittsburgh, USA; ²Leidos, USA; ³National Energy Technology Laboratory, USA. Fiber optic sensing technologies show unique relevance for energy infrastructure sensing. A non-exhaustive overview of several emerging trends within the field of optical fiber sensing technology and energy infrastructure monitoring is presented.

BM4A.6 • 18:15

Long-Term High-Temperature Wavelength Drift Trends of Type II FBGs Written in

Standard Single Mode Fibers, Robert B. Walker¹, Cyril Hnatovsky¹, Stephen J. Mihailov¹, Manny De Silva¹, Ping Lu¹, Huimin Ding¹; ¹National Research Council Canada, Canada. A comparison of Type II Fiber Bragg Grating (FBG) wavelength drift is reported for 600°C, 800°C, 900°C and 1000°C. Significant prolonged redshift was observed at 800°C, transitioning to blueshift more quickly at higher temperatures.

16:30 -- 19:00

Room: 205A

IM4B • Integrated Quantum Photonics

Presider: Nathaniel Kinsey; Virginia Commonwealth Univ., United States

IM4B.1 • 16:30 Tutorial

Development and Applications of Superconducting Single-Photon and Photon-Number

Resolving Detectors, Adriana E. Lita¹, Varun Verma¹, Dileep Reddy¹, Martin Stevens¹, Richard Mirin¹; ¹National Inst of Standards & Technology, USA. Will present an overview of superconducting single-photon detectors such as optical transition-edge sensors (TES) and superconducting nanowire single-photon detectors (SNSPD) including the operation principles and optimization of key performance metrics required for quantum information applications.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

IM4B.2 • 17:30 Invited

Epitaxial Quantum Dots for on-Chip Photonics and Long Distance Quantum Implementations, Simone Portalupi¹; ¹*IHFG, Universität Stuttgart, Germany*. We will discuss the realization of bright sources of quantum light at telecommunication wavelength via strain engineering of In(Ga)As quantum dots. Furthermore, using GaAs-based system we will show most recent achievements in photonic integrated circuits.

IM4B.3 • 18:00 Invited

Visible Light to NIR Integrated Photonics for Atomic and Quantum Applications, Daniel J. Blumenthal¹; ¹*University of California Santa Barbara, USA*. Visible light photonic integration will enable compact atomic and quantum experiments with the potential to bring about improved reliability, performance, and scaling for sensing, clocks, and compute. We discuss progress using the ultra-low loss silicon nitride platform to integrate precision lasers and photonics and higher level functions for cold and thermal neutral atom and trapped ion based systems.

IM4B.4 • 18:30 Invited

Quantum Noise Limited Homodyne Detectors Integrated Into Silicon Photonics for Quantum Technologies, Jonathan Matthews¹; ¹*University of Bristol, United Kingdom*. It is critical for any emerging technology that a path to scalable manufacture at volume is essential for impact outside the lab. Integrated quantum photonics has already revolutionized the way in which photonic quantum technology can be researched and developed. Silicon photonics offers high component density as well as CMOS compatibility for monolithic integration of photonic devices with electronics. In this talk we will cover how a particular key component for quantum technologies — quantum noise limited homodyne detectors — can be implemented in silicon photonics and how this integration can bring about improvement in key device performance metrics. Applications of these detectors include continuous variables quantum computing and communications, and detection of squeezed light. We will show how miniaturizing these quantum light detectors shrinks total detector footprint right down to 10s of micrometres squared (including the core electronic and photonic components), opens the way to scalable implementation of high numbers of detectors for a single piece of technology, and we will show how implementing the detectors with electronic-photonic integration increases speed performance beyond what is achievable without integration. Time permitting, we will show how we are using such detectors to demonstrate chip-scale sensor nodes for distributed quantum sensing.”

16:30 -- 18:30

Room: 205B

IM4G • Passive Photonic Devices

Presider: Daniele Melati; CNRS, France

IM4G.1 • 16:30 Invited

Scalable Surface Gratings for Efficient Fiber-Chip and Free-Space Optical Coupling, Radovan Korcek¹, William Fraser^{2,3}, Sara Salhi⁴, Xiaochen Xin², Quentin Wilmart⁵, David Medina⁴, Samson Edmond⁴, Thalía Domínguez Bucio⁶, Frederic Gardes⁶, Winnie N. Ye², Jens H. Schmid³, Pavel Cheben³, Daniele Melati⁴, Laurent Vivien⁴, Carlos Ramos⁴, Daniel Benedikovic¹; ¹*University of Zilina, Slovakia*; ²*Carleton university, Canada*; ³*National Research Council, Canada*; ⁴*Centre de Nanosciences et de Nanotechnologies, CNRS & Université Paris-Saclay, France*; ⁵*CEA LETI, France*; ⁶*Optoelectronic Research Centre, University of*

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Southampton, United Kingdom. Efficient optical input/output interfaces between photonic chips and fibers or free-space ports are indispensable building blocks for a wide range of applications. Here, we present our recent progress in the development of silicon nitride grating couplers and compact silicon antennas.

IM4G.2 • 17:00

Open-Source Finite Element Models for the Design of Topology Optimized Blazed Grating Under Conical Incidence, Simon Ans^{1,2}, Frédéric Zamkotsian², Guillaume Demesy¹; ¹*Institut Fresnel, France*; ²*Laboratoire d'Astrophysique de Marseille, France*. We present a suite of open-source template models based on the finite element for the topology optimization of nanophotonic structures under conical incidence, with applications to the broadband optimization of nanostructured blazed gratings.

IM4G.3 • 17:15

Highly Efficient and Compact non-Uniform Waveguide Grating Antenna for Beam Steering Application, Diksha Maurya¹; ¹*Indian Institute of Technology (ISM) Dha, India*. We proposed a highly efficient non-uniform waveguide grating antenna for beam steering in the C band. The waveguide grating antenna is optimized using the genetic algorithm to achieve high diffraction efficiency and compactness.

IM4G.4 • 17:30

Tilted Subwavelength Grating Assisted Directional Coupler Based Wavelength Division Demultiplexer, Rajarshi Guchhait¹, Ravi Roushan Kumar¹, Devendra Chack¹; ¹*Indian Institute of Technology, India*. We propose a compact dual-wavelength multiplexer using a tilted SWG-assisted Directional Coupler (DC). The device is 28.1 μm long, with insertion loss (IL) less than 0.96 dB and an extinction ratio exceeding 16.5 dB.

IM4G.5 • 17:45

Optical Delay in Subwavelength Grating Waveguides Operating Near the Bandgap, Luyao Xie¹, Lawrence R. Chen¹; ¹*McGill University, Canada*. We explore the dispersive characteristics of subwavelength grating (SWG) waveguides near the bandgap and demonstrate how they can provide additional flexibility for providing optical time delays.

IM4G.6 • 18:00 Invited

Recent Advances on Silicon-Based Metamaterial Grating Couplers, Winnie N. Ye¹; ¹*Carleton University, Canada*. Grating couplers have emerged as crucial elements for photonic integrated circuits due to their ability to provide flexible light-coupling and enable wafer-scale testing. We present our recent advances in grating coupler designs using subwavelength metamaterials.

16:30 -- 18:30

Room: 205C

JM4D • Radiative Cooling III (Joint SOLED/NOMA)

Presider: Jyotirmoy Mandal; Princeton University, United States

JM4D.1 • 16:30 Invited

Passive Radiative Cooling Film (PRCF) and Applications of PRCF, Timothy Hebrink¹; ¹*3M Company, USA*. Nano-layered and nano-particle technologies have been innovated into easy to

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

apply passive cooling films and have demonstrated significant cooling of roof surfaces by both reflecting solar energy and radiating heat away from the roof surfaces.

JM4D.2 • 17:00 Invited

Polymer-Based Multi-Functional Micro-Photonic Metamaterials for Radiative Cooling, Gan Huang¹, Bryce S. Richards¹; ¹*Karlsruher Institut für Technologie, Germany*. Polymer-based micro-photonic multi-functional materials from small-scale to large-scale, wherein micro-photonic structures are used to enable multi-function of high-emissivity, self-cleaning, and also high transparency.

JM4D.3 • 17:30

Passive Radiative Cooling With High-Mass Roof and Temperature-Driven Ventilation, Remy Fortin¹, Jyotirmoy Mandal², Salmaan Craig¹; ¹*McGill University, Canada*; ²*Princeton University, USA*. A reduced-scale model building uses a daytime radiative cooling coating ($\alpha_{\text{solar}} \sim 0.97$, $\epsilon_{\text{LWIR}} \sim 0.94$) on a high-mass, uninsulated roof without a convection guard to produce stable indoor temperatures and temperature-driven ventilation.

JM4D.4 • 17:45

Daytime Radiative Cooling Near the Equator, Jaesuk Hwang^{1,2}; ¹*The Centre for Quantum Technologies, Singapore*; ²*National University of Singapore, Singapore*. Daytime radiative cooling is challenging in the equatorial tropics, where the weather is extremely humid and cloudy. With a high degree of thermal insulation, radiative cooling can be exploited for thermal management and energy generation.

JM4D.5 • 18:00 Invited

Polymer Metasurface Radiative Cooling Film From Lab to Fab, Baohua Jia¹, Han Lin¹, Keng-Te Li¹; ¹*Royal Melbourne Institute of Technology, Australia*. We demonstrate a thin film polymer metasurface radiative cooling film enabled by periodically arranged three-dimensional (3D) trench-like structures manufactured by a roll-to-roll printing method. It exhibits superior spectral breadth, selectivity and diurnal cooling performance.

16:30 -- 18:30

Room: 206B

NpM4E • Integrated and Nonlinear Micro-Optics

Presider: Ksenia Dolgaleva; University of Ottawa, Canada

NpM4E.1 • 16:30 Invited

Optical Nonlinearities in 2D-Material-Augmented Waveguides: Enhancement Versus Counteraction, Nathalie Vermeulen¹; ¹*Vrije Universiteit Brussel, Belgium*. Combining two-dimensional materials with waveguides can lead to an enhanced nonlinear-optical performance. However, in some cases, the combined material platforms counteract each other. In this talk, I will illustrate both scenarios with two concrete examples.

NpM4E.2 • 17:00

Nonlinear Characterization of InGaAsP/InP Platform in the C-Band Telecom Range, Gabriel Flizikowski¹, Lais F. dos Santos¹, Ozan W. Oner¹, Athulya Thulaseedharan¹, Ehsan Mobini¹, Kaustubh Vyas¹, Fatemeh M. Karimi¹, Kashif M. Awan¹, Daniel Espinosa¹, Ksenia Dolgaleva¹; ¹*University of Ottawa, Canada*. We conducted four-wave mixing and nonlinear

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

absorption experiments in InGaAsP/InP waveguides of different geometries in the telecom C-band. Further, we measured the nonlinear refractive index to be $n_2=1.9 \times 10^{-13} \text{ cm}^2 \text{ W}$.

NpM4E.3 • 17:15

A Broadband Chip-Based Gallium Phosphide Traveling-Wave Parametric Amplifier, Alberto Nardi^{1,2}, Nikolai Kuznetsov², Alisa Davydova², Mikhail Churaev², Johann Riemensberger^{3,2}, Paul Seidler¹, Tobias Kippenberg²; ¹*IBM Research Europe, Zurich, Switzerland*; ²*Swiss Federal Institute of Technology Lausanne (EPFL), Switzerland*; ³*Norwegian University of Science and Technology, Norway*. We present a gallium phosphide optical traveling-wave parametric amplifier on a chip, achieving up to 35 dB of parametric gain, corresponding to a fiber-to-fiber net gain exceeding 10 dB over a bandwidth of 140 nm.

NpM4E.4 • 17:45

Kerr Switch Versus Nonlinear Thermal Effects in Whispering Gallery Resonators, Gabriele Frigenti¹, Daniele Farnesi¹, Stefano Pelli¹, Gualtiero Nunzi Conti¹, Tatyana V. Murzina², Silvia Soria¹; ¹*Ist di Fisica Applicata Nello Carrara, Italy*; ²*Department of Physics, Moscow State University, Russian Federation*. Whispering-gallery-modes resonators are effective switching devices when either coated or filled with non-linear material. We present examples of all-optical switching of hybrid WGM using polyfluorene, a methacrylate azobenzene and an acrylate derivatives.

NpM4E.5 • 18:00

Coupled Microdisk Cavities: Emission Pattern From Circular Versus Deformed Geometries, Tom S. Rodemund², Sile Nic Chormaic¹, Martina Hentschel²; ¹*Okinawa Inst of Science & Technology, Japan*; ²*Technische Universitaet Chemnitz, Germany*. In this work, we discuss mode-dependent chiral effects that appear when two dielectric limaçon cavities are coupled to each other via their optical fields, leading to far-field emission directionality.

NpM4E.6 • 18:15

Controllable Light Distributions in 1-D Microresonator Chains With Kerr-Nonlinearity, Alekhya Ghosh^{1,2}, Arghadeep Pal^{1,2}, Lewis J. Hill¹, Graeme N. Campbell^{3,1}, Toby Bi^{1,2}, Yaojing Zhang¹, Abdullah Alabbadi^{1,2}, Shuangyou Zhang¹, Pascal Del'Haye^{1,2}; ¹*Max-Planck-Inst Physik des Lichts, Germany*; ²*Physics, Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany*; ³*Physics, University of Strathclyde, United Kingdom*. We demonstrate the control of optical power patterns in one dimensional (1D) photonic lattices of coupled resonators via the Kerr effect. This can advance photonic integrated circuits, especially for light steering, and optical computing.

16:30 -- 18:30

Room: 2101

NeM4C • Advanced Core Networks

Presider: Marco Ruffini; Trinity College Dublin

NeM4C.1 • 16:30 Tutorial

Artificial Intelligence and Machine Learning in Optical Networking [Tutorial], Christine Tremblay¹; ¹*École de technologie supérieure, Canada*. In this tutorial, we explore various applications of artificial intelligence (AI) and machine learning (ML) methods aimed at improving

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

the performance, operations, and reliability of optical networks, as well as simplifying their management.

NeM4C.2 • 17:30 Invited

The Future of Global Coherent Optical Transmission Systems for Submarine and Space Applications, Hidenori Takahashi¹, Shota Ishimura¹, Takehiro Tsuritani¹; ¹*KDDI Research, Inc., Japan*. This paper reviews recent usage of digital coherent transmission technologies in submarine cable and optical satellite communications. The digital coherent technologies play important role to support the capacity demands.

NeM4C.3 • 18:00 Invited

Integrated Optical Communication and Distributed Sensing Systems, Chao Lu^{1,2}, Jingchuan Wang¹, Shaoyi Chen², Yaxi Yan¹, Tianrui Li², Liwang Lu¹, Yichang Wu², Wenjin Huang², Alan Pak Tao Lau¹, Zhaohui Li²; ¹*Hong Kong Polytechnic University, Hong Kong*; ²*School of Electronic and Information Technology, Sun Yat-sen University, China*. Recent work on integrating sensing functions into optical communication systems is described. The developed techniques are expected to enhance the reliability of optical networks and enable a range of applications in marine and urban environments.

16:30 -- 18:30

Room: 2104

SpM4H • Next Generation PON

Presider: Jun-ichi Kani; NTT Access Service Systems Laboratories, Japan

SpM4H.1 • 16:30 Invited

IM-DD for Next Generation PON Networks, Will the Trend Continue?, Vincent Houtsma¹, Dora van Veen¹; ¹*Nokia Bell Labs, USA*. We evaluate the status of current PONs and assess if the trend for next generation PONs will continue

SpM4H.2 • 17:00

Non-Orthogonal Multiplexing Access (NOMA)-Based Dense Passive Optical Networks Considering Fairness Under Differential Split Ratios, Zixian Wei¹, Jinsong Zhang¹, Weijia Li¹, Charles St-Arnault¹, Santiago Bernal¹, Mostafa Khalil¹, Ramón Gutiérrez-Castrejón^{1,2}, Lawrence R. Chen¹, David V. Plant¹; ¹*McGill University, Canada*; ²*Institute of Engineering, Universidad Nacional Autonoma de Mexico UNAM, Mexico*. We demonstrate a NOMA-based 200-G coherent PON for two access users with a differential split ratio, where the fairness and performance of far-end users with 256 SR are increased by adjusting the power allocation factor.

SpM4H.3 • 17:15

Spatial Diversity in Non-Orthogonal Multiplexing Access (NOMA)-Based Coherent Optical Transmission Under Differential Path Loss, Zixian Wei¹, Jinsong Zhang¹, Weijia Li¹, Charles St-Arnault¹, Santiago Bernal¹, Ramón Gutiérrez-Castrejón^{1,2}, Lawrence R. Chen¹, David V. Plant¹; ¹*McGill University, Canada*; ²*Institute of Engineering, Universidad Nacional Autonoma de Mexico UNAM, Mexico*. We first demonstrate a MISO coherent optical transmission based on joint NOMA and spatial diversity, where maximal ratio combining brings 1.764-dB diversity gain for two high-path-loss 120-km and 140-km branches.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

SpM4H.4 • 17:30 Invited

Cost Effective TFDM Coherent PON Enabled by Remote Optical Carrier Delivery and Optical Injection Locking, Haipeng Zhang¹, Zhensheng Jia¹, Karthik Choutagunta¹, Luis Alberto Campos¹, Curtis Knittle¹; ¹*CableLabs, USA*. We demonstrate a TFDM coherent PON with low-cost ECL-free ONU, enabled by remote optical carrier delivery through injection locking. Experimental demonstration of 50 km transmission shows similar performance to regular ECL-based systems.

SpM4H.5 • 18:00

Rate-Flexible Coherent TFDM PON With Transparent Digital Signal Processing Aided by Residual Carrier, Ziheng Zhang¹, Yixiao Zhu¹, Guangying Yang¹, Lina Man¹, Gengming Lin¹, Qunbi Zhuge¹, Weisheng Hu¹; ¹*Shanghai Jiao Tong University, China*. We leverage modulator finite extinction ratio-induced residual carrier for transparent digital signal processing in coherent time-frequency-division-multiplexing PON. We experimentally demonstrate flexible data rates from 100-Gb/s to 300-Gb/s after 20-km SSMF transmission with a 112-ns preamble.

16:30 -- 18:30

Room: 207

SoM4F • Nonlinear Fiber Optics

Presider: Alexander Heidt, Switzerland

SoM4F.1 • 16:30 Tutorial

Machine Learning for Nonlinear Fiber Optics, Goëry Genty¹; ¹*Tampere University, Finland*. In this tutorial, we will discuss how the techniques of machine learning techniques can be leveraged for the analysis, prediction, and control of nonlinear propagation dynamics in optical fiber systems.

SoM4F.2 • 17:30 Invited

Fiber Light Sources for Nonlinear Imaging Applications, Cassia Corso¹, Tigran Mansuryan², Alessandro Tonello², Lukasz Zinkiewicz^{1,3}, Bartosz Fabjanowicz¹, Mateusz Pielach¹, Agnieszka Jamrozik¹, Tomasz Kardas⁴, Yago Arosa², Piotr Wasylczyk³, Vincent Couderc², Yuriy Stepanenko¹, Katarzyna Krupa¹; ¹*Institute of Physical Chemistry PAS, Poland*; ²*Université de Limoges, XLIM UMR CNRS 7252, France*; ³*Faculty of Physics, University of Warsaw, Poland*; ⁴*Fluence Sp. z o.o., Poland*. We discuss new ideas for developing fiber light sources for nonlinear imaging. We demonstrate high-power SC-based laser tunable within fingerprint region, as well as new methods of tuning FWM sidebands.

SoM4F.3 • 18:00

Strong Reduction of Frequency-Comb Noise in All-Normal Dispersion Supercontinuum, Benoît Sierro¹, Sandro Camenzind², Benjamin Willenberg², Alexander Nussbaum-Lapping², Anupamaa Rampur¹, Ursula Keller², Christopher R. Phillips², Alexander M. Heidt¹; ¹*Institute of Applied Physics, University of Bern, Switzerland*; ²*Department of Physics, Institute for Quantum Electronics, ETH Zurich, Switzerland*. Experimental observations show that broadening frequency-combs in all-normal dispersion fibers can reduce noise by up to 20 dB. Our new numerical model propagates pulse trains generated from real noise measurements and replicates this accurately.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

SoM4F.4 • 18:15

Scaling of Stimulated Raman Scattering and Molecular Modulation in Hollow Anti-Resonant Fibers, Pau Arcos¹, Arturo Mena¹, María Sánchez-Hernández¹, Amaia Berganza², Begoña Garcia-Ramiro², Joseba Zubia⁴, David Novoa^{4,3}; ¹*Communications Engineering, University of the Basque Country, Spain*; ²*Applied Mathematics, University of the Basque Country, Spain*; ³*IKERBASQUE, Basque Foundation for Science, Spain*; ⁴*Communications Engineering and EHU Quantum Center, University of the Basque Country, Spain*. We report a scaling methodology for Raman molecular modulation dynamics in gas-filled anti-resonant fibers. The dephasing-gain length ratio allows complex nonlinear propagation dynamics to be reproduced with high fidelity under very different input conditions.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Tuesday, 30 July

10:00 -- 12:00

Room 2000B

JTu1A • Joint Plenary Posters Session

JTu1A.1

Channel Prediction and Phase Correction in a Vertical FSO Link Using Recurrent Neural Network, Souvik Sen¹, Pritam Paul¹, Rik Chattopadhyay¹; ¹*Electronics and Telecommunication, Indian Institute of Engineering Science and Technology, Shibpur, India*. We report prediction and pre-compensation of phase aberration of data in FSO link from ground to a HAP using machine learning and GS Algorithm. Probe beam is used on demand to self-correct the prediction algorithm.

JTu1A.2

Optical Studies of Rare Earth Nano Metal Oxides Prepared Using D-Glycine as Fuel in Solution Combustion Synthesis, Ashish R. Tanna¹, Dhara Maheta¹, B S. Madhukar²; ¹*RK University, India*; ²*Department of Chemistry, JSS Science and Technology University, India*. The structural and optical studies have been used for nano rare earth oxides. The refractive indices modeling has been applied for CeO₂, La₂O₃, & ThO₂ where 3.10, 4.84 & 5.76 eVs values for band gaps.

JTu1A.3

TDBC Microstructures Made by Local Photo-Bleaching in J-Aggregate Thin Organic Layers for Photonics Applications, Komlan Segbéya Gadedjisso-Tossou¹, Antoine Bard², Clementine Symonds², Jean-Michel Benoit², Joel Bellessa², Alban Gassenq²; ¹*University of Lomé, Togo*; ²*Institut Lumière Matière, France*. TDBC layers are very interesting for photonics applications due to their huge oscillator strength, narrow absorption, local photo-bleaching, and low-cost fabrication. These intrinsic properties need to be investigated to fully exploit his high potential.

JTu1A.4

Optical Manipulation of Microdroplets for Precise Imaging and Manipulation of Nanostructures, Xixi Chen¹; ¹*Institute of Nanophotonics, Jinan University, China*. The presentation centers on optical manipulation research for precise microdroplet formation/dissolution, shaping, and repositioning, enabling accurate nanostructure imaging and manipulation. It covers the physical mechanisms and fabrication methods of droplets using artificial and natural biological materials.

JTu1A.5

Analyzing Near-Field Enhancement in Plasmonic Nanocylinder-Based SPASER Systems, Amine Jaouadi¹, Ahmed Mahjoub², Montacer Dridi³; ¹*ECE - Paris School of Engineering, LyRIDS - ECE Research Center, France*; ²*Space Science Institute, USA*; ³*SCIQ, ESIEA, France*. We explore here the enhancement of electromagnetic fields near to a periodic array of gold nanoparticles. The obtained SPASER demonstrates a remarkable promise, boasting energy conversion efficiencies of up to 80%, underscoring its potential as a highly efficient nano-scale energy source.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

JTu1A.6

Epsilon Near Zero Metasurfaces (ENZ) at Visible Wavelengths, Iman Alhamdan¹; ¹*school of physics and Astronomy, United Kingdom*. Here we showcase a multilayer structure metasurface incorporating an epsilon near zero (ENZ) material. The simulation results prove the tendency of this structure to operate with high efficiency and tunability in the visible range.

JTu1A.7

Durable Fiber Bragg Gratings With Thickened Polyimide Coatings for High Sensitivity Humidity Sensing, Stephen J. Mihailov¹, Huimin Ding¹, Robert B. Walker¹, Katherine Szabo¹, Cyril Hnatovsky¹, Abdullah Rahnema¹, Ping Lu¹, Manny De Silva¹; ¹*National Research Council Canada, Canada*. Fiber Bragg Grating (FBG) relative humidity (RH) sensors were written in 50 μm diameter fibers through the polyimide coating. Thickening the polyimide resulted in the highest sensitivity reported for polyimide coated FBG RH sensors.

JTu1A.8

Specificity of NY-ESO-1 Antibody Detection Using TFBG Plasmonic Sensor, Linyao Tan², Hang Qu², Xiaoyong Chen³, Yi-Wei Xu⁴, Patrice M egret¹, Christophe Caucheteur¹, Xuehao Hu¹; ¹*Universite de Mons, Belgium*; ²*Shantou University, China*; ³*Dongguan University of Technology, China*; ⁴*Cancer Hospital of Shantou University Medical College, China*. Autoantibodies against New York esophageal squamous cell cancer 1 (NY-ESO-1) are essential for diagnosing esophageal cancer. This study introduces a surface plasmonic tilted fiber Bragg grating (TFBG) biosensor designed to detect NY-ESO-1 antibody and investigate its specificity.

JTu1A.9

Factors Influencing the Behaviour of FBG Sensors for Temperature Measurements, Tommaso Carles^{1,2}, Patrice M egret¹; ¹*University of Mons, Belgium*; ²*Environmental and Industrial Flow, Von Karman Institute for Fluid Dynamics, Belgium*. Investigating Fiber Bragg Gratings (FBG) behavior in varied fluid environments for temperature measurements. Analysis includes secondary effects and liquid metal influence on peak spectrum.

JTu1A.10

Fluoride Long Period Grating Developed Using Filament Fusion Splicer, Antreas Theodosiou^{2,1}, Yauhen Baravets², Kirill Grebnev³, Maria Chernysheva³, Pavel Honzatko², Pavel Peterka²; ¹*Lumoscribe Ltd., Cyprus*; ²*Institute of Photonics and Electronics, Czechia*; ³*Leibniz Institute of Photonic Technology, Germany*. We report on the first fabrication of long period grating in ZBLAN fiber using filament fusion splicer. The grating has been designed to operate at 1900-nm range. The sensitivity of the grating with respect to the axial tension has been investigated.

JTu1A.11

Laser-Induced Fabrication of Micro-Optics on Bioresorbable Calcium Phosphate Glass, Devanarayanan M. Menon¹, Nadia G. Boetti², Davide Janner¹; ¹*Politecnico di Torino - DISAT, Italy*; ²*LINKS Foundation, Italy*. We present a flexible and precise technique to obtain microoptics in a bioresorbable phosphate glass by laser processing. Tuning the laser parameters we obtained different micro-optic shapes from hyperbolic to parabolic lenses and diffraction gratings.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

JTu1A.12

Reference Wavelength Comb Based on High Order Fiber Bragg Grating for FBG Optical Interrogators, Romain Cotillard¹, Nicolas Roussel¹; ¹CEA, France. We use a High Order Fiber Bragg Grating as an all-fiber reference wavelength comb for a FBG measurement system based on a tunable VCSEL source.

JTu1A.13

Data Analysis Algorithm to Evaluate Temperature Tendency From Locomotives Diesel Engines, Gabriel Martelli¹, Eduardo H. Dureck¹, Felipe Mezzadri¹, Cicero Martelli¹, Jean Carlos Cardozo da Silva¹; ¹Federal University of Technology-Paraná, Brazil. The proposed algorithm interprets locomotive engine oil temperature data via the FBG sensor system, conducting a thorough analysis, reducing false alarms, and enhancing operational safety and precision compared to previous methods.

JTu1A.14

High-Temperature Wavelength Drift Comparison of Type II FBGs Written in Different Types of Single Mode Fiber, Robert B. Walker¹, Cyril Hnatovsky¹, Stephen J. Mihailov¹, Manny De Silva¹, Ping Lu¹, Huimin Ding¹; ¹National Research Council Canada, Canada. Type II Fiber Bragg Grating (FBG) wavelength drift is reported and compared for gratings in single mode fibers, having different diameter and dopant configurations. Some fibers respond with less drift than others.

JTu1A.15

Large Arrays of Low Loss Type II FBGs Written With a Phase Mask Through a Polyimide Coating, Robert B. Walker¹, Cyril Hnatovsky¹, Stephen J. Mihailov¹, Ping Lu¹, Manny De Silva¹, Huimin Ding¹; ¹National Research Council Canada, Canada. Practical quasi-distributed sensing with Fiber Bragg Gratings (FBGs) depends on the repeatable fabrication of numerous, mechanically robust, low loss FBGs. This work reports an array of 1008 FBGs written with a phase mask through polyimide.

JTu1A.16

Femtosecond Laser Direct Writing of Polarization-Controllable DBR Fiber Lasers for Harsh Environmental Sensing, Xizhen Xu¹, Jun He¹, Runxiao Chen¹, Yiping Wang¹; ¹Shenzhen University, China. We propose the fabrication of polarization-controllable DBR FLs by using a slit beam shaping femtosecond laser point-by-point technology. Experimental results show the fabricated DBR FL can withstand a high temperature up to 800 degree.

JTu1A.17

Withdrawn

JTu1A.18

A Free Space Optical Link Model for C-Band Data and Power Transmission, Idriss A. Ali¹, Paige Wilson¹, Meghan N. Beattie¹, Ryan Hogan³, Narmada Rajaram¹, Ross Cheriton², Ahmad Atieh³, Karin Hinzer¹; ¹University of Ottawa, Canada; ²National Research Council, Canada; ³Optiwave Systems Inc., Canada. We have modelled a free space optical (FSO) channel transmitting data and power at 1550 and 1520 nm, respectively, under various meteorological conditions. Quadratic Phase-Shift Keying is predicted to have the longest viable FSO range.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

JTu1A.19

All-Optical Switch and Logic Gates Using Phase Asymmetries in High-Q Resonators, Arghadeep Pal^{1,2}, Alekhya Ghosh^{1,2}, Shuangyou zhang¹, Lewis J. Hill¹, Toby Bi^{1,2}, Pascal Del'Haye^{1,2}; ¹*Max-Planck-Inst Physik des Lichts, Germany*; ²*Physics, Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany*. We demonstrate an optical switch and propose designs of photonic logic gates exploiting the Kerr-effect induced enhancement of phase asymmetries. This will be of immense importance in integrated all-optical computing.

JTu1A.20

Room-Temperature Sputtered Silicon Nitride for Soliton Microcombs, Shuangyou zhang¹, Toby Bi^{1,2}, Irina Harder¹, Olga Ohletz¹, Florentina Gannott¹, Alexander Gumann¹, Eduard Butzen¹, Yaojing Zhang¹, Lewis J. Hill¹, Pascal Del'Haye¹; ¹*Max-Planck-Inst Physik des Lichts, Germany*; ²*Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany*. We present ultralow-loss, high-thickness silicon nitride photonic circuits fabricated using room-temperature sputtering. After 800 °C annealing, we achieve propagation losses of 3.5 dB/m, enabling ring resonators with optical quality factors exceeding 10 million.

JTu1A.21

Silicon Nitride Based Planar Lightwave Circuits for Coherent Optics, Jong-Hoi Kim¹, Sang-Ho Park¹, Young-Tak Han¹, Seo-Young Lee¹, Honghwi Park¹, Dong-Hun Lee¹, Shin-Mo An¹, Won-Seok Han¹, Jang-Uk Shin¹; ¹*Electronics and Telecom Research Inst, Korea (the Republic of)*. We present planar lightwave circuits based on silicon nitride waveguides platform to realize monolithic integration of polarization beam splitters, optical 90-degree hybrids, variable optical attenuators, and micro-ring resonators for coherent detection.

JTu1A.22

Dispersion Engineering in Silicon Nitride Ring Resonators via Partial Sidewall Modulation, Masoud Kheyri^{1,2}, Shuangyou zhang¹, Toby Bi^{1,2}, Arghadeep Pal^{1,2}, Hao Zhang^{1,4}, Yaojing Zhang^{1,3}, Abdullah Alabbadi^{1,2}, Haochen Yan^{1,2}, Alekhya Ghosh^{1,2}, Lewis J. Hill¹, Pablo Bianucci⁵, Eduard Butzen¹, Florentina Gannott¹, Alexander Gumann¹, Irina Harder¹, Olga Ohletz¹, Pascal Del'Haye¹; ¹*Max Planck institute for the science of light, Germany*; ²*Department of Physics, Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany*; ³*The Chinese University of Hong Kong, China*; ⁴*National Key Laboratory of Microwave Photonics, Nanjing University of Aeronautics and Astronautics, China*; ⁵*Department of Physics, Concordia University, Canada*. We propose a method for dispersion engineering in silicon nitride microresonators via mode splitting. Mode splitting is induced by partially modulating the resonator inner-sidewall at four different segments. We report dispersion improvement over a range 100 nm.

JTu1A.23

Time-Domain Analysis of a Reflective Kerr Microcomb, Jean-Michel Vallée¹, Wei Shi¹; ¹*Centre d'optique, photonique et laser (COPL), Laval university, Canada*. We introduce a structure based on a reflective nonlinear microring for microcomb generation. By leveraging finite difference time-domain methods in our simulations, we offer a novel approach to microcomb development.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

JTu1A.24

Integrated Photon Pairs Source Based on Counter-Propagating Spontaneous Four Wave Mixing in a Silicon Nitride Chip, Gerardo Rodríguez Becerra¹; ¹*Photonics, Centro de Investigaciones en Óptica A.C., Mexico*. We report the design of an on-chip integrated photon pair source based on Counter Propagating Spontaneous Four Wave Mixing (CP-SFWM) capable of generate separable photon pairs with narrow bandwidth.

JTu1A.25

Robust Structural Design and Fabrication of High-Efficiency Functional Metasurfaces, Po-Jui Chen¹, Chia-Wei Lu², Chin-Chuan Wu¹, Wen-Chin Hsieh¹, Chung-Chih Wu^{1,2}; ¹*Graduate Institute of Electronics Engineering, National Taiwan University, Taiwan*; ²*Graduate Institute of Photonics and Optoelectronics, National Taiwan University, Taiwan*. Incomplete etching of nanostructures would reduce transmission of metasurfaces by 10%-20%. By over-etching the nanostructures, transmission can reach a value similar to or even higher than the nanostructures being precisely etched to the substrate surface.

JTu1A.26

Numerical Estimation of a Propagation Loss Evaluation Using an Asymmetric Mach-Zehnder Interferometer, Kodai Sato¹, Hiroshi Fukuda^{1,2}; ¹*Chitose Institute of Science and Technology, Japan*; ²*Silicon Research Center, Japan*. This study numerically estimated the effectiveness of an evaluation method using an asymmetric Mach-Zehnder interferometer and an embedded photodiode and revealed that this technique is applicable for propagation loss characterization in a practical range.

JTu1A.27

A Rib Waveguide Design for Reduction of Leakage Loss of TM Modes, Mir Hammadi¹, Matthew Barrett¹, Muhammad Z. Alam¹; ¹*Queens University, Canada*. We propose a rib waveguide design to address the leakage radiation inherent to silicon rib waveguides. The guide provides good tolerance to fabrication imperfections and flexibility in independent control of orthogonal polarizations.

JTu1A.28

A Switchable Polarization Converter on an Indium Phosphide Membrane, Sander Reniers¹, Jos van der Tol¹, Kevin Williams¹, Yuqing Jiao¹; ¹*TU/e, Netherlands*. We present a switchable polarization converter on an indium phosphide membrane, using a simple fabrication process. A polarization extinction ratio above 19 dB and insertion loss below 0.5 dB are demonstrated in simulation.

JTu1A.29

Higher Order Mode Analysis in Frequency Selective Negative Curvature Fibers, Muhammad Zain Siddiqui², Ahmet E. Akosman¹, Mustafa Ordu²; ¹*Roger Williams University, USA*; ²*UNAM - Institute of Materials Science and Nanotechnology, Turkey*. A novel negative curvature hollow-core fiber is numerically designed capable of filtering specific frequencies. The six-tube silica fiber strongly favors fundamental mode transmission over higher order modes despite uneven positioning of cladding elements.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

JTu1A.30

Exploring the Impact of Gap Compensation Tubes on Enhancing the Performance of Anti-Resonant Fibers, Zhaoyang Zhang¹, Yuemei Li¹, Ziyang Xiao², Yao Guo¹, Zheng Liu¹, Haobo Guo¹; ¹*Beijing University of Posts and Telecomm, China*; ²*Information and Communications branch, Jiangxi Electric Power Company, China*. By introducing Gap Compensation tubes, we have reduced the LP₀₁ mode loss to 3.487×10^{-6} dB/km, setting a new record in anti-resonant fibers. This paper will delve into the effects of Gap Compensation tubes.

JTu1A.31

Over Two-Octave Supercontinuum Generation for OAM_{19,1} Mode in Air-Core Ring Fiber, Xiaoke Wu¹, Pengfei Wang¹, Jian Yang², Yuanpeng Liu², Zhongqi Pan³, Yang Yue¹; ¹*Xi'an Jiaotong University, China*; ²*Nankai University, China*; ³*University of Louisiana at Lafayette, USA*. An air-core ring fiber with flat and near-zero dispersion is designed and simulated for broadband OAM_{19,1} supercontinuum generation. A 2410-nm supercontinuum is formed from 770 nm to 3180 nm at -40 dB, covering more than 2-octave bandwidth.

JTu1A.32

Tailoring Strut Thicknesses for Selective THz Negative Curvature Fiber Sensors, Ethan Howard¹, Julia Ward¹, Ethan Neidt¹, Riley Como¹, Ahmet E. Akosman¹; ¹*Roger Williams University, USA*. A cascaded hollow-core negative curvature fiber design is proposed for selective THz sensing. Numerical investigations suggest significant spectral shift up to 0.1 THz in relative sensitivities, enabling differentiation between analytes at certain design frequencies.

JTu1A.33

Hyperbolic Metamaterial Enhancement Based High Sensitivity Side-Polished Fiber SPR Sensors, Shiqi hu¹, Yunhan Luo¹, Yaofei Chen¹, Lei Chen¹, Gui-shi Liu¹, Zhe Chen¹; ¹*Jinan University, China*. Hyperbolic metamaterials are novel materials that excite the surface plasmon resonance (SPR) by their unique hyperbolic dispersion properties. Here, we design a composite-based fiber HMM-SPR sensor and apply it to magnetic field and temperature sensing. It provides a new research avenue for the application of high-sensitivity fiber sensors.

JTu1A.34

Acceleration of Optomechanical Droplets, Gordon Robb¹, Josh Walker¹, Gian-Luca Oppo¹, Thorsten Ackemann¹; ¹*University of Strathclyde, United Kingdom*. We describe a scheme for acceleration sensing using stable optomechanical droplets formed when a Bose–Einstein Condensate is illuminated by a far off-resonant optical pump field and by its retroreflection from a feedback mirror.

JTu1A.35

Standing-Wave Patterns Visualization on a Toroidal Microresonator, Haochen Yan^{1,2}, Alekhya Gosh^{1,2}, Arghadeep Pal^{1,2}, Hao Zhang¹, Toby Bi^{1,2}, George Ghalanos¹, Shuangyou zhang¹, Lewis J. Hill¹, Yaojing Zhang¹, Yongyong Zhuang^{1,3}, Jolly Xavier^{1,4}, Pascal Del'Haye^{1,2}; ¹*Max-Planck Institute, Science of Light, Germany*; ²*Physics, Friedrich Alexander University Erlangen-Nuremberg, Germany*; ³*Electronic Materials Research Laboratory, Xi'an Jiaotong University, China*; ⁴*SeNSE, Indian Institute of Technology, India*. The standing wave patterns generated on a microresonator by bidirectional pumping can be directly visualized by a near-infrared camera. We quantitatively analyze the scattered light intensity and apply the pattern for sub-wavelength accuracy distance measurement.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

JTu1A.36

Bound State Soliton Rain Generation in Femtosecond Fiber Laser Using Mxene Saturable Absorber, Kwani Lee¹; ¹*Korea Institute of Science & Technology, Korea (the Republic of)*. This study highlights the potential of DMSO-Ti₃C₂T_x MXene in enhancing the stability of ultrafast fiber lasers, opening avenues for exploring soliton rains in nonlinear optical dynamics.

JTu1A.37

Dispersion Engineering of Integrated Gallium Phosphide Fabry-Pérot Resonators, Alberto Nardi^{1,2}, Thomas Karg¹, Alisa Davydova², Johann Riemensberger^{3,2}, Tobias Kippenberg², Paul Seidler¹; ¹*IBM Research Europe, Zurich, Switzerland*; ²*Swiss Federal Institute of Technology Lausanne (EPFL), Switzerland*; ³*Norwegian University of Science and Technology, Norway*. We present a protocol to engineer the dispersion of integrated Fabry-Pérot resonators made of gallium phosphide by taking into account the group index of the light propagating into integrated chirped mirrors, enabling exotic dispersion profiles.

JTu1A.38

Real-Time Coupling Induced Linear and Nonlinear Light Dynamics in Coupled Microresonators, Arghadeep Pal^{1,2}, Alekhya Ghosh^{1,2}, Shuangyou Zhang¹, Lewis J. Hill¹, Haochen Yan^{1,2}, Hao Zhang^{1,3}, Toby Bi^{1,2}, Abdullah Alabbadi^{1,2}, Pascal Del'Haye^{1,2}; ¹*Max-Planck-Inst Physik des Lichts, Germany*; ²*Physics, Friedrich Alexander University, Germany*; ³*National Key Laboratory of Microwave Photonics, Nanjing University of Aeronautics and Astronautics, China*. We study the linear and nonlinear light interactions in coupled resonators. In our experiments, we can control the coupling gap between two resonators instantaneously to investigate symmetry breaking of hybridized counterpropagating modes.

JTu1A.39

Graphdiyne Oxide as a Promising Candidate for Nonlinear Optical Switching Applications, Leiming Wu¹; ¹*Guangdong University of technology, China*. Graphene oxide demonstrates a significant nonlinear optical response with light intensity dependence upon stimulation by light waves. Taking advantage of this characteristic, graphyne oxide shows important application value in optical switching.

JTu1A.40

Pulse Formation in the Stokes Region Through Dispersion Tailoring in Silica Microresonators, Ryo Otake¹, Riku Imamura¹, Shun Fuji¹, Takasumi Tanabe¹; ¹*Keio University, Japan*. We show numerically that a pulse with a peak power of ~200W is formed in the Stokes region in a silica microresonator when careful dispersion tailoring is performed.

JTu1A.41

Enhancing Solar Cell Efficiency With Tunable Transverse Modal Response to Natural Light of a Linear Chiral Thin Film Array, Monish Chatterjee¹, Akram Muntaser¹; ¹*University of Dayton, USA*. We propose the design of a chiral thin film resonator array aligned horizontally illuminated by a collimated linear *p*-polarized plane wave. This analysis aims to optimize the transmission of light; such a tuned high intensity transmission may be utilized to maximize the conversion efficiency of solar cells with tunable bandgaps.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

JTu1A.42

Tight Focusing of Obstructed Radially-Azimuthally Polarized Higher Order Poincare Sphere Beams, Sushanta K. Pal¹, Leslie Rusch¹; ¹*Université Laval, Canada*. We study the intensity landscapes of obstructed radially-azimuthally polarized higher-order Poincare sphere beams under tight focusing conditions. These optical fields can be used to achieve strong longitudinal and transverse components at the focal plane.

JTu1A.43

Engineering of Vanadium Dioxide for Reconfigurable Optics by Combining Metal Doping and Defect Engineering, Jin-Woo Cho¹, Jonathan King¹, Dung Quach², Martin Hafermann³, Karla Paz³, Hongyan Mei¹, Shenwei Yin¹, Tanuj Kumar¹, Joseph Andrade², Colin Hessel², Carsten Ronning³, David Woolf², Mikhail A. Kats¹; ¹*University of Wisconsin-Madison, USA*; ²*Physical Sciences Inc., USA*; ³*University Jena, Germany*. We demonstrate that metal doping and defect engineering can work in tandem to modify phase-transition temperature and hysteresis of vanadium dioxide films, enabling new opportunities for reconfigurable infrared optics.

JTu1A.44

A Template-Matching-Based Algorithm for Optical Microscope Image Stitching, Vikesh S. Bhadouria¹, You-rim Park¹, Joo-beom Eom¹; ¹*Department of Biomedical Science, Dankook University, Korea (the Republic of)*. This paper proposes a template-based method to enhance image-stitching in whole slide microscopy, tackling stage misalignment and flat field issues, improving accuracy and efficiency despite mechanical instability.

JTu1A.45

Characterization of Zinc Sulfide Waveguides for Nonlinear Photonics, Antoine Lemoine¹, Antoine Létoublon¹, Alex Naïm¹, Thomas Batte¹, Charles Cornet¹, Yannick Dumeige¹, Christophe Levallois¹, Yoan Léger¹; ¹*Univ Rennes, INSA Rennes, CNRS, Institut FOTON - UMR 6082, F-35000 Rennes, France, France*. This paper presents the advantages of zinc sulfide (ZnS) for second order nonlinear photonics through structural and optical characterizations of ZnS thin films and waveguides.

JTu1A.46

Withdrawn

JTu1A.47

Analysis of the Optical Behavior of Diffuse Reflectance in Polycrystalline Yttrium-Iron Garnet Synthesized by Different Methods and its Effect to Estimate Eg by Tauc Plot, Lis Tamayo-Rivera¹, Anette López-Sierra¹, Diana Salvador-García¹, Joel E. Valdivieso-Villegas¹, María del Pilar Gutiérrez-Amador¹, Ariadna Sánchez-Castillo¹; ¹*Autonomous University of Hidalgo State, Mexico*. Optical characterization of colored samples allowed us to observe clear differences in amplitude and position of reflectivity bands around the fundamental absorption energy-edge; so, different criteria to estimate Eg by Tauc plot are discussed

JTu1A.48

Withdrawn

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

JTu1A.49

Defect Superlattice Optical Solitons Supported by Centrosymmetric Photonic Superlattice, Draupath Umesh¹, Gaurang Potdar¹, Aavishkar Katti¹; ¹*MIT World Peace University, India*. Optical spatial gap solitons are analyzed in centrosymmetric photorefractive media with an embedded defect superlattice for the first time. Double and multi hump solitons existence and characteristics are studied across both band gaps.

JTu1A.50

Exploring Bistable Properties in AgCl-Ag Multilayer Nonlinear Structures for Blood Serum Analysis, Iuliia Riabenko¹, Konstantin Beloshenko¹; ¹*V. N. Karazin Kharkiv National University, Ukraine*. Silver transfer to interference minima in AgCl-Ag films creates periodic structures through interaction with the waveguide TE₀-mode. This phenomenon enables the recording of Raman spectra of biological samples, potentially enhancing low-intensity signal registration.

14:00 -- 16:00

Room: 206A

BTu2A • Symposium on Optical Fiber Sensors for Extreme Environments II

Presider: Tobias Habisreuther; Leibniz Institute of Photonic Tech., Germany

BTu2A.1 • 14:00 Invited

Sensing in Extreme High Temperature Environments With Regenerated Fiber Bragg Gratings, Johannes Roths¹, Qiang Bian², Andrea Stadler¹, Fabian Buchfellner¹, Alexander Roehrl¹; ¹*Photonics Laboratory, Munich University of Applied Sciences, China*; ²*College of Meteorology and Oceanography, National University of Defense Technology, China*. In recent years, multipoint temperature sensors based on regenerated fibre Bragg gratings (RFBG) have achieved a high degree of maturity. Specific sensor characteristics and applications in metallurgy, gas turbine and aircraft engine instrumentations, are reviewed.

BTu2A.2 • 14:30

Nanoscale Porous Silica Sensing Layers on Tilted FBGs Made by Flame Spray Pyrolysis, Zayne Ramotar¹, Qianzhu Li², Jacques Albert², Reza Kholghy¹, Hubert Jean-Ruel²; ¹*Department of Mechanical and Aerospace Engineering, Carleton University, Canada*; ²*Department of Electronics, Carleton University, Canada*. Nanoscale coatings of pure porous silica are deposited on optical fibers in a few minutes and atmospheric conditions. A tilted FBG in the fiber enables measurements of the coating and of its sensing properties.

BTu2A.3 • 14:45

Point-by-Point Femtosecond Fiber Bragg Gratings Behavior at High Temperatures, Matilde Sosa^{1,2}, Maxime Cavillon², Thomas Blanchet¹, Matthieu Lancry², Guillaume Laffont¹; ¹*Université Paris-Saclay, CEA List, France*; ²*Institut de Chimie Moléculaire et des Matériaux d'Orsay (ICMMO/SP2M/MAP), Université Paris-Saclay, France*. Void-based fiber Bragg gratings were studied from quantitative phase microscopy and spectral measurements, under isochronal annealing experiments up to 1250°C. We reveal a link between the micro-void deformation and the Bragg reflectivity degradation at high temperatures.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

BTu2A.4 • 15:00

High Temperature Resistance fs-FBGs in Large Diameter Optical Fibers, Karima Chah¹, Damien Kinet^{1,2}, Corentin Guyot², Christophe Caucheteur¹; ¹UMONS / *Faculté Polytechnique de Mons, Belgium*; ²B-SENS, *Belgium*. We subject Fs-FBGs in silica optical fibers of 125, 200 and 400 μm diameters to high temperature cycling. Post-annealing tensile tests confirm the higher mechanical resistance of large diameter optical fibers compared to standard ones

BTu2A.5 • 15:15

Withdrawn

BTu2A.6 • 15:30 Invited

Interferometric Inscription of Sapphire Fiber Bragg Gratings and Temperature Diagnosis Using These Gratings in Applications Above 1400°C, Tobias Habisreuther¹, Kerstin Schröder¹, Rene Eisermann², Adrian Lorenz¹, Stephan Krenek²; ¹*Leibniz Institute of Photonic Tech., Germany*; ²*Physikalisch-Technische Bundesanstalt, Germany*. The paper reports the inscription of fiber Bragg gratings by 400nm femtosecond laser pulses using an interferometer setup. High temperatures probes were assembled, calibrated using temperature fixed-points and applied in industrial processes at $T > 1400^\circ\text{C}$.

14:00 -- 16:00

Room: 205A

ITu2B • Optical Communications and Computing

Presider: Nathalie Vermeulen; Vrije Universiteit Brussel, Belgium

ITu2B.1 • 14:00 Invited

Ultrafast Membrane Lasers With Optical Feedback for Optical Interconnects and Neuromorphic Computing, Pandelis Diamantopoulos¹, Takuro Fujii¹, Suguru Yamaoka¹, Hidetaka Nishi¹, Koji Takeda¹, Shinji Matsuo¹; ¹*NTT Device Technology Labs, Japan*. We present the first 16-channel membrane laser array exhibiting photon-photon resonance, achieving 50-GHz bandwidths and < 130 fJ/bit energies, for 1.6 Tbps transceivers. Additionally, we have showcased ultra-fast and energy-efficient spiking dynamics for neuromorphic applications.

ITu2B.2 • 14:30

BER Measurement in Data Transmission With a Double-Pass SOA, Hiroya Sakumoto^{1,2}, Yi Wang², Desalegn W. feyisa², Sander Reniers², Ripalta Stabile², Kevin Williams², Yuqing Jiao²; ¹*Department of Electrical Engineering and Information Systems, The University of Tokyo, Japan*; ²*Eindhoven Hendrik Casimir Institute, Eindhoven University of Technology, Netherlands*. We demonstrate transmitting NRZ-OOK PRBS31 signal at 25 Gb/s with a double-pass SOA, showing a 61% gain boosted with the same electrical power consumption while keeping almost the same BER as a conventional single-pass SOA.

ITu2B.3 • 14:45

Flexible Opto-Electronic Logical Gate Circuit Comprising Waveguide-Based Interferometer and Photodetector, Koichi Takiguchi¹, Hironori Nishihara¹; ¹*Department of Electrical and Electronic Engineering, Ritsumeikan University, Japan*. We report an opto-electronic logical gate with flexible operation, which consists of an integrated-optic symmetric

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Mach-Zehnder interferometer and a balanced photodetector. We show Boolean AND and NAND computations of 40 Gbit/s signals with the gate.

ITu2B.4 • 15:00

Compact TE-Pass Polarizer Covering Full Communication Band With Ultra-High Extinction Ratio in O-Band and C+L Band, Zhuoran Wang¹; ¹*McGill University, Canada*. We design and simulate a hybrid plasmonic waveguide TE-pass polarizer operating from 1260 - 1675 nm with an extinction ratio >18 dB, and exceeding 35 dB in the O band and C+L band with bandwidths of 50 nm and 120 nm, respectively.

ITu2B.5 • 15:15

Proposal for a 500 GHz Silicon Photonic Modulator, Conglin Sun^{1,2}, Minkyu Kim¹, Pol Van Dorpe^{1,2}, Francky Catthoor^{1,3}, Mikael Mazur⁴, Joris Van Campenhout¹, Christian Haffner¹, Dennis Lin¹; ¹*Interuniversity Microelectronics Centre, Belgium*; ²*Department of Physics and Astronomy Research unit Quantum Solid-State Physics, KU Leuven, Belgium*; ³*Department of Electrical Engineering (ESAT), KU Leuven, Belgium*; ⁴*Nokia Bell Labs, USA*. We propose a silicon modulator concept utilizing organic electro-optical Pockels materials in a vertical gap configuration to reach 500 GHz bandwidth. Simulations show that optical losses can be ~0.5 dB/~0.2 dB for a 500 GHz/200 GHz design with an V_{π} of 13.3 V/5.3 V respectively

ITu2B.6 • 15:30 Invited

Non-Reciprocal Materials for Photonic in-Memory Computing, Nathan Youngblood¹, Paolo Pintus^{2,3}, Mario Dumont³, Vivswan Shah¹, Toshiya Murai⁵, Yuya Shoji⁴, Duanni Huang³, John Bowers³; ¹*University of Pittsburgh, USA*; ²*Physics, University of Cagliari, Italy*; ³*ECE, University of California, Santa Barbara, USA*; ⁴*Electrical and Electronic Engineering, Tokyo Institute of Technology, Japan*; ⁵*Platform Photonics Research Center, National Institute of Advanced Industrial Science and Technology (AIST), Japan*. Non-reciprocal platforms can offer several key advantages for scalable and efficient photonic computing. In this talk, I will present our recent experimental work validating the use of non-reciprocal materials to implement high-endurance memory for photonic computing.

14:00 -- 16:00

Room: 206B

NpTu2E • Applications of Nonlinear Optics

Presider: Sife Nic Chormaic; Okinawa Inst of Science & Technology, Japan

NpTu2E.1 • 14:00 Invited

NLO Microscopy, Randy A. Bartels¹; ¹*Morgridge Institute & U of Wisconsin, USA*. Abstract not available.

NpTu2E.2 • 14:30 Invited

Miniaturized Nonlinear Micro-Endoscopes, Dylan Septier², Eloïse Lefebvre^{1,2}, Gaëlle Brévalle-Wasilewski², Naveen Gajendra Kumar², Yong Jian Wang², Attila Kaszas², Hervé Rigneault³, Alexandre Kudlinski¹; ¹*Univ Lille 1 Laboratoire PhLAM, France*; ²*Lightcore Technologies, France*; ³*Institut Fresnel, Aix Marseille Univ, Centrale Med, France*. We report miniaturized and flexible micro-endoscopes based on double-clad negative curvature hollow core fibers. They allow to perform multiphoton, second and third harmonic and coherent anti-Stokes Raman scattering imaging.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

NpTu2E.3 • 15:00

Study of Composite Optical Nanofibers for 2nd and 3rd Order Nonlinearities, Sylvie Lebrun¹, Abderrahim Azzoune², Maha Bouhadida¹, Théo Damp³, Laurent Divay³, Mathieu Fauvel³, Christian Larat³, Jean-Charles Beugnot⁴; ¹*Université Paris-Saclay, Institut d'Optique Graduate School, CNRS, Laboratoire Charles Fabry, France*; ²*Ecole Militaire Polytechnique, Laboratoire Systèmes Lasers, Algeria*; ³*Thales Research and Technology, France*; ⁴*Institut Femto-ST, CNRS, Université Bourgogne, Franche-Comté, France*. We present the design of composite optical nanofibers coated with different nonlinear materials for the realization of 2nd and 3rd order nonlinear effects.

NpTu2E.4 • 15:15

1-GHz Dual-Comb Supercontinuum From a Single Nonlinear Fiber Using Polarization Multiplexing, Alexander M. Heidt¹, Sandro Camenzind², Benoît Sierro¹, Anupamaa Rampur¹, Benjamin Willenberg², Ursula Keller², Christopher R. Phillips²; ¹*Universität Bern, Switzerland*; ²*Institute for Quantum Electronics, ETH Zurich, Switzerland*. We report 1-GHz watt-level ultra-low noise dual-comb supercontinuum generation in a single birefringent photonic crystal fiber using polarization multiplexing. Both combs originate from the same laser cavity and are broadened to > 450 nm bandwidth.

NpTu2E.5 • 15:45

All-Fiber Broadband mid-Infrared Supercontinuum Generation in as₂S₃-Polycarbonate Hybrid Microtaper, Md. Hosne Mobarok Shamim¹, Imtiaz Alamgir¹, Martin Rochette¹; ¹*McGill University, Canada*. We present an all-fiber supercontinuum source spanning over the spectral range of 1.2-3.9 μm from an As₂S₃-polycarbonate hybrid microtaper. This is the broadest supercontinuum obtained from a robust polymer clad hybrid microtaper.

14:00 -- 16:00

Room: 205C

NoTu2D • Emerging Imaging Techniques for Biology and Materials Science (Computational Methods)

Presider: Alon Gorodetsky; University of California Irvine, United States

NoTu2D.1 • 14:00 Tutorial

Holotomography and Artificial Intelligence: Label-Free 3D Imaging, Classification, and Inference of Live Cells, Tissues, and Organoids, YongKeun Park¹; ¹*Korea Advanced Inst of Science & Tech, Korea (the Republic of)*. Holotomography (HT) is a powerful label-free imaging technique that enables high-resolution, three-dimensional quantitative phase imaging (QPI) of live cells and organoids through the use of refractive index (RI) distributions as intrinsic imaging contrast¹⁻³. Similar to X-ray computed tomography, HT acquires multiple two-dimensional holograms of a sample at various illumination angles, from which a 3D RI distribution of the sample is reconstructed by inversely solving the wave equation.

NoTu2D.2 • 15:00 Invited

3D Inverse-Scattering in Biological Samples, Shwetadwip Chowdhury¹; ¹*University of Texas at Austin, USA*. I will cover the computational imaging frameworks that support our recent efforts for 3D imaging of multiple-scattering samples. I will specifically focus on our choice of large-scale nonlinear and nonconvex inverse-scattering techniques. Additionally, I will share recent

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

findings from applying these methods to biological samples, and I will explore ongoing challenges and prospective avenues for future research.

NoTu2D.3 • 15:30 Invited

On the use of Machine Learning for Quantifying Complex Processes, With Application to Retina Vasculature and Glaucoma Diagnostics, George Barbastathis^{1,2}, Sandip Mondal², Thiara S. Ahmed³, Bingayo Tan^{4,3}, Qihang Zhang⁵, Fabian Braeu⁴, Liangcai Cao⁵, Michael Girard⁴, Leopold Schmetterer^{4,3}, Aung Tin⁴; ¹*Massachusetts Institute of Technology, USA*; ²*Singapore-MIT Alliance for Research and Technology Centre, Singapore*; ³*Nanyang Technological University, Singapore*; ⁴*Singapore Eye Research Institute, Singapore*; ⁵*Tsinghua University, China*. We investigate the mapping between flows on complex graphs, such as retina vasculature, and spatial-temporal fluctuations in the far field. Unsupervised and supervised learning algorithms can be used to invert these maps and obtain quantitative information about connectivity and blood flow.

14:00 -- 16:00

Room: 2101

NeTu2C • SDM and Multi-band Networks

Presider: Ruben Luis; NICT, Japan

NeTu2C.1 • 14:00 Invited

Withdrawn

NeTu2C.2 • 14:30 Invited

High Capacity Optical Transmission Using Spatial Division Multiplexing, Georg Rademacher¹; ¹*Universität Stuttgart, Germany*. Space-division multiplexing offers a tremendous potential for high speed optical fiber communications systems. In this talk, we will review key components and technologies required to implement SDM transmission in multi-mode and multi-core fibers.

NeTu2C.3 • 15:00 Invited

Raman Amplifiers for Multi-Band Optical Transmission Systems, Mingming Tan¹, Pratim Hazarika¹, Dini Pratiwi¹, Wladek Forysiak¹; ¹*Aston University, United Kingdom*. We showcase effective strategies to mitigate undesirable pump-to-pump power transfer in wideband Raman amplifiers, encompassing a spectrum of up to 210nm, inclusive of E, S, C, and L bands

NeTu2C.4 • 15:30

MPI Impact in C+L+S Multiband Transmission Reach, Luis G. Cancela^{1,2}, João O. Pires^{3,2}; ¹*Iscte - Instituto Universitário de Lisboa, Portugal*; ²*Instituto de Telecomunicações, Portugal*; ³*Instituto Superior Técnico, Portugal*. Multipath interference (MPI) impact is assessed in a C+L+S multiband transmission scenario. For a typical -34 dB/span MPI, transmission reach in the L-band suffers a 20% reach penalty considering the QPSK modulation format.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

14:00 -- 16:00

Room: 2104

SpTu2H • Next Generation PON, Fronthaul and Data Center Networks

Presider: Molly Piels; OpenLight, United States

SpTu2H.1 • 14:00 Invited

Softwarization of DSP and Other Physical-Layer Functions for Future Optical Access, Sang-Yeup Kim¹; ¹*NTT Access Service Systems Laboratories, Japan*. This talk covers challenges and opportunities encountered when designing future optical access given the wide variation in service requirements. To address the challenges, we introduce a softwarized platform that offers DSP-oriented access on commodity servers.

SpTu2H.2 • 14:30

Single-Wavelength 100 Gbit/s PS-PAM8 Transmission Over Hybrid Fiber-FSO Links for Data Center Interconnect, Ahmed Galib Reza¹, Lakshmi Narayanan Venkatasubramani¹, Anil Raj Gautam¹, Liam P. Barry¹; ¹*Dublin City University, Ireland*. We demonstrate ≈ 100 Gbit/s PS-PAM8 transmission over a hybrid intra-data center interconnect link consisting of 1 km SMF and 6 m FSO channels. The BER below the HD-FEC level is achieved with a T-spaced FFE.

SpTu2H.3 • 14:45

RL-Based Digital Pre-Distortion for Drive Signals in Non-Differentiable Channel, Arash Rabieepoor¹, Leslie Rusch¹, Ming Zeng¹; ¹*Université Laval, Canada*. This paper proposes a novel digital pre-distortion, a technique based on deep reinforcement learning to combat instantaneous nonlinearities in electrical back-to-back communication systems. Simulation results demonstrate its superiority over the case without DPD in terms of bit error rate performance.

SpTu2H.4 • 15:00 Invited

Digital-Analog Mobile Fronthaul With Efficient SNR Scaling, Yixiao Zhu¹, Chenbo Zhang², Xiansong Fang², Yicheng Xu¹, Ziheng Zhang¹, Fan Zhang², Qunbi Zhuge¹, Xiaopeng Xie², Weisheng Hu¹; ¹*Shanghai Jiao Tong University, China*; ²*Peking University, China*. We overview the principle of hybrid digital-analog radio-over-fiber technique for high-fidelity and high spectral efficiency mobile fronthaul. The SNR scaling law is theoretically derived and compared with the existing radio-over-fiber techniques.

SpTu2H.5 • 15:30

Analog FFE Coefficients Optimization Using MMSE-Based LMS Algorithm in PON Context, Dylan chevalier^{1,2}, Pascal Scalart³, Gaël Simon¹, Laurent Bramerie², Michel Joindot², Jérémy Potet¹, Mathilde Gay², Philippe Chanclou¹, Monique Thual²; ¹*Orange Labs, France*; ²*Institut FOTON - UMR 6082, France*; ³*IRISA - UMR 6074, France*. A novel approach is proposed for Analog Signal Processing (ASP) in enhanced 50G-PON using analog FFE filters based on MMSE technique. Performance are validated through experimental transmission of electrical NRZ signals.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

14:00 -- 16:00

Room: 205B

STu2G • Solar Optics

Presider: Mathieu de Lafontaine; University of Ottawa, Canada

STu2G.1 • 14:00 Invited

New Limits for Light-Trapping and Ultrathin Solar Cells, Stéphane Collin^{1,2}, Maxime Giteau¹; ¹C2N CNRS, France; ²IPVF, France. We present new upper bounds for light-trapping in solar cells, and we provide an answer to the long-debated question of the best strategy for light-trapping: isotropic scattering using random texturing, or multi-resonant absorption using periodical patterning. We also discuss state-of-the-art ultrathin solar cells and prospects.

STu2G.2 • 14:30

Stationary Solar Concentrators for any Latitude, Gianpaolo Lenarduzzi¹, Tahzinul Islam¹, Thomas Cooper¹; ¹York University, Canada. We introduce a generalized source/acceptance map matching method to design non-tracking concentrators for virtually any latitude. Several practical configurations are explored. A particularly useful arrangement makes use of a pivotable front wall to achieve a seasonally-adaptive design which surpasses the static concentration limit.

STu2G.3 • 14:45

Daylighting System Using Solar Concentrator With Efficient Spectral Utilization of Sunlight, Mayank Gupta^{1,2}, Zala P. Bharatsinh¹; ¹Department of Physics, Pandit Deendayal Energy University, India; ²Indian Institute of Technology Delhi, India. Spectral profile of Sunlight is utilized efficiently with solar Fresnel lens as primary and Compound parabolic as secondary concentrator for daylighting. The results of photometric parameters of daylighting are very encouraging with improved optical efficiency.

STu2G.4 • 15:00 Invited

Title to be Announced, Andru Prescod¹; ¹ManTech International Corp., USA. Abstract not available.

STu2G.5 • 15:30

A Roadmap on Optics for Terawatt Scale Photovoltaics, Klaus Jaeger^{1,2}, Sven Burger^{2,3}, Urs Aeberhard⁴, Esther Alarcon Llado⁵, Benedikt Bläsi⁶, Bruno Ehrler⁵, Wilfried Favre⁷, Antonin Fejfar⁸, Tristan Gageot⁷, Ivan Gordon^{9,10}, Henning Helmers⁶, Oliver Höhn⁶, Olindo Isabella¹⁰, Marko Jost¹¹, Martin Ledinsky⁸, Jyotirmoy Mandal¹², Phillip Manley^{3,2}, Delfina Munoz⁷, Juan C. Ortiz Lizcano¹⁰, Ulrich W. Paetzold¹³, Aaswath P. Raman¹⁴, Hitoshi Sai¹⁵, Rebecca Saive¹⁶, Martina Schmid¹⁷, Eli Yablonovitch¹⁸, Christiane Becker¹; ¹Helmholtz-Zentrum Berlin für Materialien und Energie GmbH, Germany; ²Zuse Institute Berlin, Germany; ³JCMwave GmbH, Germany; ⁴Fluxim AG, Switzerland; ⁵AMOLF, Netherlands; ⁶Fraunhofer ISE, Germany; ⁷CEA-INES, LITEN, France; ⁸FZU - Institute of Physics of the Czech Academy of Sciences, Czechia; ⁹IMEC, Belgium; ¹⁰Delft University of Technology, Netherlands; ¹¹University of Ljubljana, Slovenia; ¹²Princeton University, USA; ¹³Karlsruhe Institute of Technology, Germany; ¹⁴University of California Los Angeles, USA; ¹⁵AIST, Japan; ¹⁶University of Twente, Netherlands; ¹⁷Universität Duisburg-Essen, Germany; ¹⁸University of California Berkeley, USA. The ongoing development of photovoltaics into terawatt scale poses a number of challenges where the optics

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

and photonics communities can contribute. An international consortium recently compiled a roadmap that elaborates on these challenges.

STu2G.6 • 15:45

Optimization of 1.65 eV $\text{Al}_{0.18}\text{Ga}_{0.82}\text{As}$ Tunnel Junctions for Monolithic III-v/Si Solar Cell, May Angelu L. Madarang^{2,1}, Rafael Jumar Chu^{2,1}, Yeonhwa Kim^{2,3}, Eunkyo Ju^{2,3}, Tsimafei Laryn^{1,2}, Won Jun Choi², Daehwan Jung^{1,2}; ¹*Nanomaterials Science and Technology, KIST School at University of Science and Technology, Korea (the Republic of)*; ²*Center for Optoelectronic Materials and Devices, Korea Institute of Science and Technology, Korea (the Republic of)*; ³*Department of Materials Science and Engineering, Korea University, Korea (the Republic of)*. We investigate thermally stable, optically transparent 1.65 eV AlGaAs tunnel junctions crucial for interconnecting tandem cells. Devices with 2x 8% Te delta-doping show five orders of magnitude improvement and thermal robustness compared to Si-doped counterparts.

14:00 -- 16:00

Room: 207

SoTu2F • Trends in Industry & Commercial Applications

Presider: Yingying Wang; Jinan University, United States

SoTu2F.1 • 14:00 Invited

Modern Industrial Processes to Bring Specialty Optical Fibre to Scale, Julien Roy¹; ¹*Coractive, Canada*. Specialty Optical Fiber demand and performance have greatly evolved in the last 25 years. In this talk we present the major challenges faced by this industry and how Coractive has embraced these market evolutions.

SoTu2F.2 • 14:30 Invited

Hollow Core Fibre: a Specialty Fibre With the Dream of Becoming Mainstream, Francesco Poletti¹; ¹*University of Southampton, United Kingdom*. Abstract not available.

SoTu2F.3 • 15:00 Invited

Polarization Maintaining Large-Mode Area Double Clad Fiber Design Strategies to Power-Scale Narrow-Line Width Industrial Lasers Beyond 2 kW, Clemence Jollivet¹; ¹*Coherent Corp, USA*. Abstract not available.

SoTu2F.4 • 15:30

Withdrawn

16:30 -- 18:30

Room: 206A

BTu3A • Glasses & Gratings in Radiative Environment

Presider: Matthieu Lancry; Universite Paris-Saclay, France

BTu3A.1 • 16:30 Invited

Femtosecond Laser Fabrication Enabled Fiber Optical Sensors for Nuclear Energy Applications: Rad-Hard Sensors, Sensor Fusion, and Machine Learning, Kevin P. Chen¹; ¹*University of Pittsburgh, USA*. Abstract note available.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

BTu3A.2 • 17:00

Behavior During in-Core Reactor Irradiation at 500°C of Regenerated and Type III Femtosecond FBGs Written in Germano-Silicate Single-Mode Fiber, Rudy Desmarchelier¹, Romain Cotillard¹, Matthieu Lancry², Stephane Breaud³, Andrei Goussarov⁴, Christophe Destouches³, Guillaume Laffont¹; ¹CEA Paris-Saclay, France; ²Université Paris-Saclay, France; ³CEA Cadarache, France; ⁴SCK-CEN, Belgium. In-core reactor irradiation of regenerated and type III femtosecond Fiber Bragg Gratings has been realized during the TESCA experiment in the SCK CEN BR2 reactor on the LIBERTY rig. FBG transducers are simultaneously sensitive to environment. In this study we have investigated a new method to discriminate temperature and radiation effects on FBGs.

BTu3A.3 • 17:15

Radiation Induced Refractive Index Change in Optical Fibres Through Rayleigh-OFDR and FBG Techniques, Jérémy Perrot¹, James Hainsworth^{1,2}, Emmanuel Marin¹, Adriana Morana¹, Youcef Ouerdane¹, Aziz Boukenter¹, Hugo Boiron³, Johan Bertrand⁴, Sylvain Girard¹; ¹Institut d'Optique Lab Hubert Curien, France; ²IRT Saint Exupery, France; ³Exail, France; ⁴ANDRA, France. This study compares radiation-induced refractive index variations in SMF-28e+ fibres using Type II femtosecond-inscribed FBGs and R-OFDR techniques. It aims to understand femtosecond laser pulse effects on fibre core composition and behaviour in harsh environment.

BTu3A.4 • 17:30

Type III Femtosecond Fiber Bragg Grating Behaviors Under X-Rays, Thomas Blanchet¹, Benjamin Sapaly¹, Romain Cotillard¹, Sylvain Magne¹, Adriana Morana², Emmanuel Marin², Sylvain Girard², Christophe Destouches³, Guillaume Laffont¹; ¹CEA Saclay - LIST - LSPM, France; ²Hubert Curien Laboratory, France; ³CEA DES Cadarache, France. We demonstrate that the behavior under X-rays at a constant dose-rate at room temperature of type III fs-void fiber Bragg gratings, inscribed with the point-by-point technique, depends on the fiber radiation response, i.e. on its core composition.

BTu3A.5 • 17:45 Invited

Withdrawn

16:30 -- 18:30

Room: 205A

ITu3B • Integrated Photonic Devices I

Presider: Christian Haffner; Interuniversity Microelectronics Center, Belgium

ITu3B.1 • 16:30 Invited

Overcoming Fundamental Noise Sources in Parametric Oscillators, Krishna Twayana¹, Fuchuan Lei^{1,2}, Victor Torres Company¹; ¹Chalmers University of Technology, Sweden; ²Northeast Normal University, China. We demonstrate self-injection locking of a parametric oscillator subject to feedback. This process results in phase noise reduction of modes below the limits imposed by fundamental noise sources in the coherent parametric wave generation.

ITu3B.2 • 17:00

Silicon Nitride Microring Resonator for Optical Frequency Comb Generation in Normal Dispersion, Rifat Nazneen¹, Odile Liboiron-Ladouceur¹; ¹McGill University, Canada. We

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

measure a SiN microring resonator with a Q-factor of 7.78×10^5 . Simulation shows that the resonator can generate six comb lines with 60 mW sources using four-wave mixing in normal dispersion.

ITu3B.3 • 17:15

Mode-Dependent Thermo-Optic Phase Shifter Using Coupled Waveguides on Silicon for 2- μ m Waveband, Taichi Muratsubaki¹, Takanori Sato¹, Kunimasa Saitoh¹; ¹*Hokkaido University, Japan*. Silicon mode-dependent thermo-optic phase shifter is proposed for the 2- μ m waveband. Appropriately designed coupled-waveguides structure provides a difference in the thermo-optic phase-shift factors between TE₀ and TE₁ modes by a factor of two.

ITu3B.4 • 17:30

InAs/GaAs Quantum dot Based Laser Arrays and Single Photon Sources, Ying Yu¹, Jaiwei Yang¹, Hancheng Zhong¹, Siyuan Yu¹; ¹*Sun Yat-Sen University, China*. We have presented low noise distributed-feedback laser arrays and low-threshold continuous-wave micro-lasers based on quantum dots (QDs), and also demonstrated a bright single photon source from a deterministically coupled single QD within monolithic Fabry-Perot microcavity.

ITu3B.5 • 17:45

Theoretical Exploration of Biosensing Using Hybrid Semiconductor Plasmonic Lasers, Shayan Saeidi¹, Pavel Cheben², Jens H. Schmid², Pierre Berini¹; ¹*University of Ottawa, Canada*; ²*National Research Council of Canada, Canada*. We present a new methodology for biosensing based on a hybrid plasmonic-semiconductor laser. We use the laser metal contact for plasmonic sensing while using an electrically pumped semiconductor heterostructure to produce TM gain.

ITu3B.6 • 18:00 Invited

Withdrawn

16:30 -- 18:30

Room: 206B

NpTu3E • Frequency Conversion

Presider: Sylvie Lebrun; *Centre National Recherche Scientifique, France*

NpTu3E.1 • 16:30 Invited

Material Engineering for Poling sub- μ m Periods in KTP, Laura Barrett¹, Cherrie Lee¹, Carlota Canalias¹; ¹*Kungliga Tekniska Hogskolan, Sweden*. We present and discuss different methods for domain engineering in KTP. These methods are based on ion-exchange, are useful for fabrication of bulk nano-domain gratings, and are compatible with waveguide implementation.

NpTu3E.2 • 17:00

Simultaneous Modal Phase- and Group Velocity Matching for Frequency Tripling and Consecutive Cascaded Wave Mixing, Artemii Tishchenko¹, Francis Berghmans¹, Tigran Baghdasaryan¹; ¹*Vrije Universiteit Brussel, Belgium*. We show numerically the feasibility to achieve doubly phase-matched cascaded four-wave mixing in high GeO₂-content microstructured optical fibers. The first stage of said cascade involves simultaneous modal phase and group velocity matching-based third harmonic generation

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

NpTu3E.3 • 17:15 Invited

Adiabatic Frequency Converter as a Single-Cycle Pulse Generator and Custom Dispersive Element, Jeffrey Moses¹; ¹*Cornell University, USA*. An adiabatic parametric frequency converter efficiently handles octave-spanning bandwidth while providing a route to tailor group delay dispersion through the quasi-phase matching grating design. Recent advancements include nearly dispersion-free frequency translation of mid-infrared single-cycle pulses.

NpTu3E.4 • 17:45

Over 2.5-Octave OAM Supercontinuum Generation in ZBLAN Ring-Core Fiber, Haoyang Ren¹, Wenpu Geng², Jian Yang², Yuetian Wang³, Zhongqi Pan⁴, Yang Yue¹; ¹*School of Information and Communications Engineering, Xi'an Jiaotong University, China*; ²*Institute of Modern Optics, Nankai University, China*; ³*Department of Optoelectronic Information Science and Engineering, School of Physics, Xi'an Jiaotong University, China*; ⁴*Department of Electrical & Computer Engineering, University of Louisiana at Lafayette, USA*. A ring-core ZBLAN fiber with low dispersion is designed and simulated for broadband OAM supercontinuum generation. A 4117-nm supercontinuum is formed from 863 nm to 4980 nm at -40 dB, covering more than 2.5-octave bandwidth.

NpTu3E.5 • 18:00

Tavis-Cummings Model for Frequency Up-Conversion in Gas-Filled Anti-Resonant Hollow-Core Fibers, Tasio Gonzalez-Raya^{1,2}, Luca Leggio^{1,3}, Arturo Mena³, David Novoa^{4,5}, Mikel Sanz^{6,7}; ¹*Basque Center for Applied Mathematics (BCAM), Spain*; ²*EHU Quantum Center, Spain*; ³*Department of Communications Engineering, University of the Basque Country (UPV/EHU), Spain*; ⁴*Department of Communications Engineering, University of the Basque Country (UPV/EHU) and EHU Quantum Center, Spain*; ⁵*IKERBASQUE, Basque Foundation for Science, Spain*; ⁶*Basque Center for Applied Mathematics (BCAM) and IKERBASQUE, Basque Foundation for Science, Spain*; ⁷*Department of Physical Chemistry, University of the Basque Country (UPV/EHU) and EHU Quantum Center, Spain*. We introduce a full quantum-mechanical description of recent experiments on correlation-preserving frequency up-conversion of single photons in gas-filled anti-resonant fibers using an N-molecule Tavis-Cummings model. Our formalism predicts preservation of entanglement under specific conditions.

NpTu3E.6 • 18:15

All-Fiber Coherent Supercontinuum in the mid-Infrared, Md. Hosne Mobarok Shamim¹, Laurent Brilland², Radwan Chahal², Johann Troles², Martin Rochette¹; ¹*McGill University, Canada*; ²*CNRS, ISCR-UMR, Université Rennes, France*. We present an all-fiber coherent supercontinuum spanning over the spectral range of 1.7- 5.0 μm from a cascade of silica, ZBLAN, and chalcogenide nonlinear fibers. The estimated average coherence across the supercontinuum is 0.83.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

16:30 -- 18:30

Room: 2101

NpTu3C • Soliton Dynamics

Presider: Silvia Soria; Inst di Fisica Applicata Nello Carrara, Italy

NpTu3C.1 • 16:30 Invited

Emergence of Laser Cavity-Solitons in Micro-Resonators, Alessia Pasquazi¹, Gian-Luca Oppo²; ¹Loughborough University, United Kingdom; ²Physics, University of Strathclyde, United Kingdom. We review our results on the modelling and observation of temporal laser cavity-solitons in a microresonator filtered fibre laser.

NpTu3C.2 • 17:00

Withdrawn

NpTu3C.3 • 17:15

Soliton Self-Generation Under Pulsed-Pumping Conditions in a Coherently Driven Passive Kerr Resonator, Matthew Macnaughtan^{1,2}, Zongda Li^{1,2}, Yiqing Xu^{1,2}, Xiaoming Wei³, Zhongmin Yang³, Stephane Coen^{1,2}, Miro Erkintalo^{1,2}, Stuart Murdoch^{1,2}; ¹The University of Auckland, New Zealand; ²The Dodd-Walls Centre for Photonic and Quantum Technologies, New Zealand; ³School of Physics and Optoelectronics, South China University of Technology, China. We unveil a novel dynamical regime in which dissipative Kerr cavity solitons spontaneously emerge in a coherent, pulse-driven passive Kerr resonator. The regime is robust against perturbations, ensuring reliable and deterministic cavity soliton generation.

NpTu3C.4 • 17:30

Demonstration of the in-Amplifier Soliton Self-Frequency Shift Optimization by Pre-Chirping, Robi Kormokar¹, Martin Rochette¹; ¹McGill University, Canada. We experimentally demonstrate that in-amplifier soliton self-frequency shift and energy conversion efficiency are maximized using a pump pulse with chirp of $C_0 \approx 0.65g_{LD}$. This result is fundamental for optimal design of SSFS based wavelength converters.

NpTu3C.5 • 17:45

Pre-Compression of High-Order Soliton for Enhanced Soliton Self-Frequency Shift, Fariha Mehjabin¹, Md Hosne Mobarok Shamim¹, Martin Rochette¹; ¹McGill University, Canada. We demonstrate the optimal design conditions of a soliton self-frequency shift (SSFS) based wavelength converter with pulse pre-compression. Sidelobes energy of a compressed high-order soliton is partially recycled to maximize SSFS and energy conversion efficiency.

NpTu3C.6 • 18:00 Invited

Dissipative Temporal Solitons in Coherently Driven Phase Modulated Cavities and in Active PT-Symmetric Dimers, Simon-Pierre Gorza¹; ¹Universite libre de Bruxelles, Belgium. We discuss recent results on the manipulation of temporal Kerr cavity solitons by external potentials and on the spontaneous formation of pulses by mode-locking in active PT-cavities. Our experimental demonstrations are performed with fiber resonators.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

16:30 -- 18:30

Room: 205C

NoTu3D • Emerging Imaging Techniques for Biology and Materials Science (Experimental Strategies)

Presider: Alon Gorodetsky; University of California Irvine, United States

NoTu3D.1 • 16:30 Invited

Quantitative Phase Microscopy With Sub-Å Measurement Accuracy for Mapping Interlayer Spacing of 2D Materials, Nansen Zhou¹, Renjie Zhou¹; ¹*Chinese University of Hong Kong, Hong Kong*. We developed quantitative phase microscopy with pm-level sensitivity. By introducing a practical phase cavity on the sample substrate, we achieved 0.1 Å measurement accuracy for mapping electronic coupling induced interlayer spacing in 2D materials.

NoTu3D.2 • 17:00 Invited

Seeing the Invisible: Meta-Optics and Phase Imaging, Ann Roberts¹; ¹*University of Melbourne, Australia*. Recent demonstrations of all-optical, object plane image processing using metasurfaces highlight their considerable potential as alternative, energy-efficient, compact, analog computing platforms. Here, their application to phase contrast imaging, including of unstained biological cells, is discussed.

NoTu3D.3 • 17:30 Invited

Advances in Quantitative Label-Free Imaging for Micro/Nano-Scale Object Analysis, Jose A. Rodrigo¹, Tatiana Alieva¹; ¹*Universidad Complutense de Madrid, Spain*. Quantitative label-free imaging provides information about the 3D shape of micro/nano-scale objects and their electric permittivity. Advantages and drawbacks of refractive index tomography and a technique based on nanoparticle manipulation by structured light are discussed.

NoTu3D.4 • 18:00 Invited

Microwave Resonances in Water-Based Objects, Yuchen Song¹, Aaron D. Slepko¹; ¹*Trent University, Canada*. We study centimeter-scale aqueous objects as high-index models for fundamental microwave dielectric resonators. We use thermal imaging, calorimetry, FEM simulations, and analytical methods to predict how/which resonances in isolated spheres couple to form intense dimer hotspots.

16:30 -- 18:30

Room: 2104

NoTu3H • Reconfigurable Materials and Devices

Presider: Richard Osgood; US Army, United States

NoTu3H.1 • 16:30 Invited

Withdrawn

NoTu3H.2 • 17:00

Optimising Liquid Crystal Alignment via Image Gradient Analysis, Zihan Feng¹, Francois Ladouceur¹, Amr A. Abed¹, Reem Almasri¹; ¹*University of New South Wales, Australia*. This paper explores the quantification of liquid crystal alignment, crucial for optimising optoelectronic devices. By using image processing techniques, we facilitate the quantification of alignment thus facilitating the identification of alignment-related issues during device fabrication.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

NoTu3H.3 • 17:15

Impact of Lattice Strain on Vanadium Dioxide Transition Temperature, Induced via Defects and Doping, Jonathan King¹, Jin-Woo Cho¹, Dung Quach², Martin Hafermann³, Karla Paz³, Hongyan Mei¹, Shenwei Yin¹, Tanuj Kumar¹, Joseph Andrade², Colin Hessel², Carsten Ronning³, David Woolf², Mikhail A. Kats¹; ¹*University of Wisconsin-Madison, USA*; ²*Physical Sciences Incorporated, USA*; ³*Friedrich Schiller University Jena, Germany*. We modify the transition temperature of vanadium dioxide films using two different methods: Ar⁺ irradiation and transition-metal-doping and observe a unified relationship between transition temperature and lattice strain, regardless of modification method or combination thereof.

NoTu3H.4 • 17:30

Hollow Plasmonic 3D Metamaterials for LDOS Enhancement and Control of Light-Matter Interaction, Margoth Córdova¹; ¹*University of Ottawa, Dept of Physics, Canada*. We proposed to explore room-temperature quantum devices operating at the single-photon level, with functionalities that can be tailored changing different parameters of a plasmonic metamaterial composed of hollow truncated nanocones that greatly enhance the photonic density of states.

NoTu3H.5 • 17:45 Postdeadline Submission

Cephalopod-Inspired Optical Living Systems With Tunable Properties, Nikhil Kaimal¹, Georgii Bogdanov¹, Alon Gorodetsky¹; ¹*University of California Irvine, USA*. Cephalopods are powerful sources of inspiration for the engineering of dynamic optical systems. We have drawn inspiration from cephalopod skin cells to engineer human cells to possess tunable transparency-changing and light scattering capabilities.

16:30 -- 18:30

Room: 205B

STu3G • Photonic Devices

Presider: Noel Giebink; University of Michigan, United States

STu3G.1 • 16:30 Invited

Colloidal Quantum Dot Laser Diodes and ASE Light Source, Victor I. Klimov¹, Valerio Pinchetti¹; ¹*Los Alamos National Laboratory, USA*. Solution-processable colloidal quantum dot lasers and amplifiers have been pursued for ease of integration with on-chip circuits. Here we discuss recent progress and present challenges in this field with focus on electrically pumped devices.

STu3G.2 • 17:00 Invited

OLEDs With Record Light Output Enable an Electrically Driven Polymer Laser, Ifor D. W. Samuel¹; ¹*University of St Andrews, United Kingdom*. Organic semiconductors have many attractive properties and are suitable as laser gain media. However, making electrically driven organic lasers has proved exceptionally challenging. This is because of the low mobility of the materials, losses due to charges, and losses due to triplets and contacts. We have explored an approach of separating charges and light generation from the gain medium to address the main challenges. This in turn brings its own challenges as it requires the development of OLEDs giving world record intensity of light output, and efficient transfer of the light generated to the laser structure. We show that such an OLED can excite lasing in a polymer distributed feedback laser. Measurements of threshold, beam and spectral narrowing will be presented.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

STu3G.3 • 17:30 Invited

3D Interconnects and III-v Semiconductor Plasma Etching for Low-Cost and High-Efficiency Photonic Devices, Mathieu de Lafontaine¹; ¹*University of Ottawa, Canada*. Photonic chips require new processes to enable transitioning to 3D interconnects. We fabricated 3D interconnects on a multijunction solar cell. We demonstrate photonic devices having areas 3 orders of magnitude smaller compared to standard chips.

STu3G.4 • 18:00 Invited

Withdrawn

STu3G.5 • 18:00 Invited

Withdrawn

16:30 -- 18:30

Room: 207

SoTu3F • Soft Glass Fibers

Presider: Solenn Cozic; Le Verre Fluore, France

SoTu3F.1 • 16:30 Invited

High Power Diode Pumped Rare Earth Doped ZBLAN Visible Fiber Lasers: Technologies, Challenges and Opportunities, Thierry Georges¹; ¹*Oxxius SA, France*. Abstract not available.

SoTu3F.2 • 17:00 Invited

Phosphate Glass Fibers for Biomedical Applications, Nadia G. Boetti¹, Jawad Talekkara Pandayil^{1,2}, Sharon Russo², Martha Segura³, Diego Pugliese², Joris Lousteau⁴, Davide Janner²; ¹*Fondazione LINKS, Italy*; ²*DISAT, Politecnico di Torino, Italy*; ³*Universitat Rovira i Virgili (URV), Spain*; ⁴*DCMC, Politecnico di Milano, Italy*. Phosphate glass optical fibers hold promise in biomedical applications, providing versatility for diagnosis, monitoring and treatment. Tailoring specific glass compositions enables biocompatibility and resorbability, facilitating implantation and unlocking potential for various promising applications.

SoTu3F.3 • 17:30

Low-Loss Single-Mode Fluoride Optical Fiber Coupler, Gebrehiwot T. Zeweldi¹, Mohsen Rezaei¹, Martin Rochette¹; ¹*McGill University, Canada*. A single-mode fluoride optical fiber coupler is demonstrated with excess loss of ≤ 0.75 dB in the spectral range of 1500-2680 nm. Surface crystallization and associated losses are reduced by processing under an argon environment.

SoTu3F.4 • 17:45

Germanate Glass: Enabling 2 μm Eye-Safe Lasers, Nadia G. Boetti¹, Martha Segura², Amiel Ishaaya³, Davide Janner⁴, Francesc Diaz², Xavier Matteos², Joris Lousteau⁵; ¹*Fondazione LINKS, Italy*; ²*Universitat Rovira i Virgili, Spain*; ³*Ben-Gurion University of the Negev, Israel*; ⁴*Politecnico di Torino, Italy*; ⁵*Politecnico di Milano, Italy*. We report on the design, fabrication, and characterization of a novel germanate glass host for efficient 2 μm eye-safe laser operation. Spectroscopic characterization and CW laser generation with Tm³⁺ and Ho³⁺ doping is investigated.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

SoTu3F.5 • 18:00

In-Line Polarization Control in Soft Glass Fibers, Md Moinul Islam Khan¹, Md Hosne Mobarok Shamim¹, Martin Rochette¹; ¹*McGill University, Canada*. We demonstrate the first proof of concept in-line polarization control using soft-glass fibers, crucial for mid-infrared applications. ZBLAN and hybrid As₂S₃ fibers exhibit high polarization extinction ratios (PER) of 20.8 dB and 19.8 dB, respectively.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Wednesday, 31 July

09:00 -- 11:00

Room: 2000A

JW1A • Plenary Session II

JW1A.1 • Plenary Submission

An Overview of Data Movement in AI/ML Systems and Their Impact on Interconnects, Ashkan Seyedi¹; ¹*NVIDIA Corporation, USA*. This talk will provide an overview of how system topologies, workloads and AI algorithms drive specific demands for system interconnects for data movement. A state-of-the-art summary of leading interconnect solutions will provide the audience with an update on how copper and optical interconnects stand to solve these problems. The second portion of the talk will focus on upcoming trends and technologies for future systems, such as classical optical and in-memory computing, as well as packaging and thermo-mechanical challenges.

JW1A.2 • Plenary Submission

A Light in Digital Darkness: Optical Wireless Communication to Connect the Unconnected, Mohamed-Slim Alouini¹; ¹*King Abdullah University of Science and Technology (KAUST), Saudi Arabia*. The transformative influence of Internet and Communication Technology (ICT) has reshaped society, touching every aspect from the economy to healthcare. As the widespread deployment of 5G continues, there is an ongoing focus on the inception of the sixth generation (6G) of wireless communication systems (WCSs). Anticipated to shape the future of connectivity in the 2030s, 6G aims to deliver unparalleled communication services to meet hyper-connectivity demands. While densely populated urban areas have traditionally been the primary beneficiaries of WCS advancements, the vision for 6G transcends city limits. Aligned with the United Nations' sustainability goals for 2030, an important aspect of 6G endeavors to democratize the benefits of ICT, fostering global connectivity sustainably. This talk delves into this particular envisioned landscape of 6G, providing insights into the future of wireless communication and guiding research efforts toward sustainable, inclusive and high-speed connectivity solutions for the future. Central to this discussion are two emerging technologies: Free Space Optics (FSO) and Non-Terrestrial Networks (NTN). These innovative solutions hold the promise of extending high-speed connectivity beyond urban hubs to underserved regions, fostering digital inclusivity and contributing to the development of remote areas. Through this exploration, we aim to convey the potential of 6G and its role in shaping a connected, sustainable future for all.

14:30 -- 16:00

Room: 206A

BW2A • BGPP Industry Session

Presider: Remco Nieuwland; Somni Solutions, The Netherlands

BW2A.1 • 14:30 Invited

High Performance Distributed Acoustic Sensing Enabled by Continuously Enhanced Backscattering Fiber, Ping Lu¹; ¹*OFS Fitel LLC, USA*. We developed a grating-based specialty single-mode fiber that is compatible with most distributed acoustic/vibrational sensing interrogators. Laboratory and field-based testing results with improved sensing performance including SNR and position accuracy will be discussed.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

BW2A.2 • 15:00 Invited

The BraggATune: the Ultimate Fiber Bragg Grating Writing Machine, Raman Kashyap¹; ¹*Photonova Inc., Canada*. We present our fiber Bragg grating tunable fabrication system. A motorized system capable of adjusting itself to write at any wavelengths, with a single phase-mask. Our system can also provide custom apodization, phase-shifts and chirp.

BW2A.3 • 15:15 Invited

Opportunities and Challenges in the Photonics Market, David Melanson¹, Julien Gagnon¹; ¹*ITF Technologies Inc., Canada*. This presentation provides a high-level view of the opportunities and challenges in the world of photonics components and sub-systems. From market demand to new applications, we will see what drives today's photonic experts.

BW2A.4 • 15:30 Invited

Fiber-Based Sensing Using Optical Frequency Domain Reflectometry (OFDR) and Distributed Bragg Gratings, Alex Tongue¹; ¹*Sensuron, USA*. Abstract not available.

BW2A.5 • 15:45

Femtosecond Laser Writing of Fiber Bragg Grating Within a PM Active Fiber for 1535 nm all-Fiber Laser Demonstration, Alain Abou Khalil¹, Valerian Freysz¹, Kevin Armengaud¹, Christophe Pierre¹, Marc Castaing¹; ¹*ALPhANOV, France*. Using femtosecond laser pulses, a highly reflective Fiber Bragg Grating was inscribed inside the core of an active PM fiber, integrated into a 1535 nm all-fiber PM laser system

14:30 -- 16:00

Room: 205A

IW2B • Neural Network Photonics

Presider: Nathaniel Kinsey; Virginia Commonwealth Univ., United States

IW2B.2 • 14:30

Photonic Neural Network and in-Situ Training in a Synthetic Frequency Dimension, Felix Gottlieb¹, Abhinav Sinha², Kai Wang¹; ¹*McGill University, Canada*; ²*Indian Institute of Science (IISc), India*. We develop a scalable photonic neural network utilizing the discrete frequency degree of freedom of light with the ability to train itself based on an in-situ backpropagation method with minimal reliance on external computers.

IW2B.3 • 14:45

Inverse-Designed 16-Channel Time-of-Flight Receiver in 45nm Silicon Photonic Process, John Rollinson¹, Robert F. Karlicek¹, Mona M. Hella¹; ¹*Rensselaer Polytechnic Institute, USA*. A 16-channel monolithic electronic-photonic receiver for flash time-of-flight LIDAR is demonstrated in 45nm silicon photonic process. Enverse-designed grating array antennas are utilized for efficient, low-noise coupling of free-space light, thus improving signal-to-noise ratio and extending sensing range of the free-space optical receiver.

IW2B.1 • 15:00 Invited

Mastering Silicon Photonics Device Design for Scalable and Robust Optical Neural Networks, Zahra Ghanaatian¹, Amin Shafiee¹, Mahdi Nikdast¹; ¹*Colorado State University, USA*. We demonstrate a design optimization approach for Mach-Zehnder Interferometers

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

(MZIs) in optical neural networks under fabrication-process variations (FPVs). Our results show increased inferencing accuracy in the network under realistic FPVs by using optimized MZIs.

IW2B.4 • 15:30

Physical Neural Networks Based on Multimode Optical Waves, Logan Wright¹; ¹*Yale University, USA*. Physical neural networks provide a way to realize neural network calculations by leveraging the controllable computations physical systems natively perform. I present an example on-chip physical neural network based on arbitrarily controllable multimode wave propagation.

14:30 -- 16:00

Room: 206B

NpW2E • Integrated Nonlinear Photonics

Presider: Silvia Soria; Inst. di Fisica Applicata Nello Carrara, Italy

NpW2E.1 • 14:30 Invited

Ultrafast Optics on Thin-Film Lithium Niobate, Mengjie Yu¹; ¹*University of Southern California, USA*. Abstract not available.

NpW2E.2 • 15:00 Invited

THz Generation Using Thin-Film LNBO, Ileana-Cristina Benea-Chelmus¹; ¹*Ecole Polytechnique Federale de Lausanne, Switzerland*. Abstract not available.

NpW2E.3 • 15:30

Impact of Nonlinear Losses on the Performance of InGaAsP-on-Insulator Ring Resonators, Athulya Thulaseedharan¹, Laís Fujii dos Santos¹, Tara Moradi², Connor Kupchak², Ksenia Dolgaleva¹; ¹*University of Ottawa, Canada*; ²*Carleton University, Canada*. We study the impact of nonlinear losses such as two-photon absorption (TPA) and free-carrier absorption (FCA) on frequency comb generation in ring resonators made of InGaAsP (In_{0.7}Ga_{0.18}As_{0.3}P on SiO₂ (InGaAsP-on-insulator, or InGaAsP-OI) platform.

14:30 -- 16:00

Room: 205C

NoW2D • Metasurfaces and Inverse Design

Presider: Lynda Busse; US Naval Research Laboratory, United States

NoW2D.1 • 14:30 Invited

Physical-Model-Based Wave Control With Reverberation-Nonlocal Programmable Metasurfaces, Philipp del Hougne¹; ¹*IETR, CNRS, IETR - Univ Rennes, France*. Chaotic-cavity-backed programmable metasurfaces experience significant coupling between meta-atoms (i.e., non-locality) due to reverberation. We frugally calibrate accurate physical models mapping metasurface configuration to scattered fields and discuss how they enable previously inaccessible wave-control regimes.

NoW2D.2 • 15:00 Invited

All-Glass Metasurfaces for High Power Lasers Optics Using a Self-Organizing Approach, Eyal Feigenbaum¹; ¹*Lawrence Livermore National Laboratory, USA*. We present a scalable method for producing all-glass meta-surfaces using a self-organizing approach, resulting in high

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

mechanical stability and laser-induced damage durability. We will present process advances enabling formation of lenses, antireflection surfaces, and waveplates.

NoW2D.3 • 15:30

Inverse-Designed Metasurface for Multidimensional Spatial State Reconstruction, Yuming Niu¹, Kai Wang¹; ¹*McGill University, Canada*. We report inverse-designed nonlocal metasurfaces for transforming multidimensional states of light represented in the Hermite-Gaussian basis into optimally designed spatial states, where a simple imaging can accurately extract the full multidimensional state including amplitude, phase, and coherence.

NoW2D.4 • 15:45

Inverse Design of Photonic Systems, Benjamin MacLellan^{1,2}, Piotr Roztock^{1,2}, Julie Belleville¹, Luis Romero Cortés^{1,3}, Kaleb Ruscitti¹, Bennet Fischer¹, José Azaña¹, Roberto Morandotti¹; ¹*Institut National de la Recherche Scientifique, Canada*; ²*Ki3 Photonics Technologies, Canada*; ³*Universitat Politècnica de València, Spain*. We present a framework for identifying the optimal topologies and operational parameters of photonic systems. Leveraging automatic differentiation and topology search, it facilitates the discovery of physically-feasible designs for applications like waveform generation and sensing.

14:30 -- 16:00

Room: 2101

NeW2C • ML and AI informed Networks

Presider: Lena Wosinska; Chalmers Tekniska Högskola, Sweden

NeW2C.1 • 14:30 Tutorial

AI-Driven Optical Network Automation: Opportunities and Challenges, Carlos Natalino¹; ¹*Chalmers Tekniska Högskola, Sweden*. Abstract not available

NeW2C.2 • 15:30 Invited

Withdrawn

14:30 -- 16:30

Room: 2104

SpW2H • Quantum Communication and Computing

Presider: Stephan Pachnicke; Christian-Albrechts Universität zu Kiel, Germany

SpW2H.1 • 14:30 Invited

Time-Multiplexed Programmable Continuous-Variable Photonic Quantum Computing, Shuntaro Takeda¹; ¹*The University of Tokyo, Japan*. Abstract not available.

SpW2H.2 • 15:00

Digital Equalization Techniques for Continuous Variable Quantum Key Distribution, Utku Akin^{1,2}, Jonas Berl^{1,3}, Tobias Fehenberger¹, Norbert Hanik²; ¹*Adva Network Security GmbH, Germany*; ²*Institute for Communications Engineering, Technical University of Munich, Germany*; ³*Communications Engineering Lab, Karlsruhe Institute of Technology, Germany*. We experimentally investigate two digital equalization schemes for ISI mitigation in CV-QKD. The

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

proposed static zero-forcing method shows similar performance to an adaptive equalizer at a significantly reduced DSP complexity.

SpW2H.3 • 15:15

Rate-Adaptive Protograph-Based Raptor-Like LDPC Code for Continuous-Variable Quantum Key Distribution, Erdem E. Cil¹, Laurent Schmalen¹; ¹*CEL, Karlsruhe Institute of Technology, Germany*. We propose a new type-based protograph raptor-like LDPC code for rate-adaptive information reconciliation in CV-QKD systems. It offers robust error-correction performance across a wide range of rates while simplifying the code design process.

SpW2H.4 • 15:30 Invited

Integrating Quantum Communications in Telecom Infrastructures, Andrew Shields¹, Tina Muller¹; ¹*Toshiba Research Europe Ltd, United Kingdom*. Abstract not available.

SpW2H.5 • 16:00

Spectral Efficient Physical Layer Security by Using Coherent All-Optical Processing and Multichannel Obfuscation, Yarden Yalievich^{1,2}, Eyal Wohlgemuth^{1,2}, Dan Sadot^{1,2}; ¹*Ben-Gurion University, Israel*; ²*CyberRidge, Israel*. A spectral-efficient physical layer security system is introduced, which enhances capacity by transmitting multiple channels over shared spectrum. Demonstrating that the success probability of an attack is extremely low and further decreases with the inclusion of more channels.

SpW2H.6 • 16:15

Quantum-Inspired Encryption Using Displacement Operators in Coherent Optical Communications, Mostafa Khalil¹, Adrian Chan², Lawrence R. Chen¹, David V. Plant¹, Randy Kuang³; ¹*McGill University, Canada*; ²*Synopsys Inc., Canada*; ³*Quantropi Inc., Canada*. We demonstrate experimental results of a physical layer encryption technique based on displacement operators in coherent optical transmission systems. The system is tested over 80 km of SSMF using 4/16/32QAM at 56 GBd.

14:30 -- 16:00

Room: 205B

SW2G • Thin-Film Applications

Presider: Stéphane Collin; CNRS, France

SW2G.1 • 14:30 Invited

Noble Transparent Conductive Oxide (TCO) Materials for Solar Cells: a Viable Alternative to Conventional TCOs, Takashi Koida¹; ¹*Natl Inst of Adv Industrial Sci & Tech, Japan*. Two transparent conductive oxide (TCO) films of interest in the solar cell field are highlighted: high-mobility In₂O₃-based TCOs currently in production lines and the indium-free, highly conductive amorphous SnO₂ films discovered recently in the laboratory.

SW2G.2 • 15:00 Invited

Contact Engineering for Organic Photovoltaics, Canek Fuentes-Hernandez^{1,2}, Felipe A. Larrain², Vladimir Kolesov², Yi-Chien Chang^{2,4}, Minwoo Nam^{2,5}, Tzu-Yen Huang³, Michael Toney³, Bernard Kippelen²; ¹*Department of Electrical and Computer Engineering, Northeastern University, USA*; ²*Center for Organic Photonics and Electronics (COPE), School of Electrical and Computer Engineering, Georgia Institute of Technology, USA*; ³*Stanford Synchrotron*

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Radiation Light Source (SSRL), SLAC National Accelerator Laboratory, USA; ⁴Faculty of Engineering and Sciences, Universidad Adolfo Ibanez, Chile; ⁵Department of Electronic Engineering, Keimyung University, Korea (the Republic of). We present an overview of solution-based electrical p-doping of organic semiconductors using polyoxometalates yielding stable exponentially decaying depth-profiles into the bulk and other highly desirable properties for the realization of organic photovoltaics.

SW2G.3 • 15:30 Invited

Material Design of Solution-Processed Semiconductors for Wearable Applications, Xiwen Gong¹; ¹*University of Michigan, USA*. I will discuss new material design strategies that impart mechanical deformability in solution-processed optoelectronic materials and their applications in wearable and healthcare technologies.

14:30 -- 16:00

Room: 207

SoW2F • Fiber Design and Modelling

Presider: Peter Dragic; Univ of Illinois at Urbana-Champaign, United States

SoW2F.1 • 14:30

Modeling Transverse Mode Instability Experiment Using Phase-Matched Model, Josh Young^{1,2}, Zhihao Hu¹, Curtis Menyuk², Jonathan Hu¹; ¹*Baylor University, USA*; ²*University of Maryland Baltimore County, USA*. We compare the results of experiment and phase-matched model for transverse mode instability (TMI). The simulation accurately identifies the threshold of TMI, which shows strong potential for future design optimizations.

SoW2F.2 • 14:45

Performance Trade-Offs Spanning O-C-L-Bands in Antiresonant Fiber Designs, Rania A. Abouelela¹, Sophie LaRoche¹, Leslie Rusch¹; ¹*ECE Department-Center for Optics, Photonics and Lasers (COPL), Laval University, Canada*. We examine antiresonant nodeless fiber (ANF) geometries with good C-band compromises between single-mode operation and loss. We identify one geometry offering good trade-offs throughout the antiresonance window, particularly in the telecommunications O-band and L-band.

SoW2F.3 • 15:00

Comparative Modelling and Design Aspects of PANDA PM Fiber, Natasha Vukovic¹, Christophe A. Codemard², Michalis N. Zervas¹; ¹*University of Southampton, United Kingdom*; ²*TRUMF Lasers UK Ltd, United Kingdom*. Using finite-element-method, we assess birefringence in PANDA PM-fiber, comparing it with several analytical expressions proposed in the literature. Our study shows agreement with one expression, while two others show varied discrepancies, indicating different underlying causes.

SoW2F.4 • 15:15

A New Simplified Bend Loss Model for Polarizing Optical Fibers, Andy Gillooly¹, Mark Hill¹, Jose Maria Alvarez De Con¹; ¹*Fibercore Limited, United Kingdom*. A simplified numerical bend loss model is presented to predict the performance of polarizing optical fibers in the spectral domain. Theoretical calculation are compared to experimental results and shown to give a close approximation.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

SoW2F.5 • 15:30

Design of a Few-Mode Fiber With Radial Anisotropy for Improved Modal Efficiency, Asma Mimouni¹, Younès Messaddeq², Bora Ung¹; ¹*Ecole de Technologie Supérieure, Canada*; ²*Centre for Optics, Photonics and Lasers (COPL), Université Laval, Canada*. A novel few-mode fiber design with radial anisotropy is proposed. The fiber improves the degeneracy of TE₀₁ and TM₀₁ modes and is a potential solution for expanding data channels and improving modal efficiency.

SoW2F.6 • 15:45

Trench-Assisted Seven-Core Non-Zero Dispersion Shifted Ring Fiber for OAM Modes, Yuxiang Huang¹, Yuanpeng Liu², Wenqian Zhao¹, Zhongqi Pan³, Yang Yue¹; ¹*Xi'an Jiaotong University, China*; ²*Nankai University, China*; ³*University of Louisiana at Lafayette, USA*. A trench-assisted seven-core non-zero dispersion shifted ring fiber is designed to support 98 OAM modes, which can maintain <-40 dB crosstalk across the C-band. Large effective mode area and low differential mode delay are also achieved.

16:30 -- 18:30

Room: 206A

BW3A • Mid-IR Glasses and Optical Materials

Presider: Daniel Richter; Institute of Applied Physics (Germany), Germany

BW3A.1 • 16:30 Invited

Fibre Bragg Gratings Inscription in Fluoride Fibers and Their Application for Spectrometry and Thermal Sensors, Maria Chernysheva¹; ¹*Ultrafast Fibre Lasers, Leibniz Institute of Photonic Technology, Germany*. In my presentation, I will demonstrate that FBGs inscribed in fluoride-based fibres can serve as sensitive thermal sensors down to cryogenic conditions as well as exhibit unique scattering patterns, facilitating tailored light emission designs.

BW3A.2 • 17:00

Thick Femtosecond-Inscribed Silica VBGs for mid-IR Spectral Filtering Applications, Lauris Talbot², Malte P. Siems¹, Daniel Richter¹, Nicolas David³, Sébastien Blais-Ouellette³, Stefan Nolte^{1,4}, Martin Bernier²; ¹*Institute of Applied Physics, Friedrich Schiller University Jena, Germany*; ²*Centre d'Optique, Photonique et Laser (COPL), Université Laval, Canada*; ³*Photon Etc., Canada*; ⁴*Fraunhofer Institute for Applied Optics and Precision Engineering IOF, Center of Excellence in Photonics, Germany*. We report on the femtosecond inscription of a 3 mm thick silica volume Bragg grating. By combining it with a mid-IR supercontinuum laser, a narrowband light source tunable from 2.8 to 4.3 μm is obtained.

BW3A.3 • 17:15

Fs-Written FBGs in InF₃ Fibers Using the Scanning Phase Mask Technique for Mid-IR All-Fiber Laser, Tommy T. Boilard¹, Réal Vallée¹, Martin Bernier¹; ¹*COPL, Université Laval, Canada*. Highly reflective FBGs are written in InF₃ fibers using the scanning phase mask technique. Their application as reflectors in an Ho³⁺:InF₃ all-fiber laser cavity emitting at 3920 nm is also presented.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

BW3A.4 • 17:30

Thermal Stability of Fiber Bragg Gratings Fabricated in Fluoride Glass Optical Fibers via Femtosecond Laser Direct-Inscription, Alex Fuerbach¹, Toney T. Fernandez^{2,1}, Luyi Xu¹; ¹*Macquarie University, Australia*; ²*University of South Australia, Australia*. We report on the annealing properties of fiber Bragg gratings (FBGs) that have been inscribed into fluorozirconate (ZBLAN) and fluoroindate (InF₃) optical fibers via the femtosecond laser direct write technique.

BW3A.5 • 17:45

Bragg Grating Reflectors Inscribed in Polypropylene Lightpipes, Vasilis Sarakatsianos¹, Ivan Chapalo¹, Eleni Grantzioti¹, Theodoros Manouras^{1,2}, Maria Vamvakaki^{1,2}, Maria Konstantaki¹, Stavros Pissadakis¹; ¹*FORTH-IESL, Greece*; ²*Department of Materials Science and Technology, University of Crete, Greece*. Bragg grating reflectors are inscribed and characterized in toluene loaded, polypropylene lightpipes using 248nm, excimer laser radiation. Refractive index changes of the order of $\sim 6.6 \times 10^{-4}$ are introduced in the polymer matrix, through single photon-absorption.

BW3A.6 • 18:00 Invited

A Type-1 Fluoride Glass Optical Waveguide Laser for mid Infrared Integrated Optics, Toney T. Fernandez¹, Dale Otten¹, Simon Gross², Michael J. Withford², Alex Fuerbach², David Lancaster¹; ¹*University of South Australia, Australia*; ²*Macquarie University, Australia*. Reporting the long-anticipated type-1 waveguide laser in erbium-doped fluoride glass. The laser cavity is formed using a butt-coupled HR-mirror and a fiber Bragg grating output coupler positioned at both ends of a 10 mm-long glass waveguide.

16:30 -- 18:30

Room: 205A

IW3B • Photonic Integrated Devices II

Presider: Judith Su; Univ of Arizona, Coll of Opt Sciences, United States

IW3B.1 • 16:30 Invited

Title to be Announced, Remus Nicolaescu¹; ¹*Pointcloud Inc., Switzerland*. Abstract not available.

IW3B.2 • 17:00

Direct Laser Writing of a Compact 1x4 Splitter for Multi-Core Optical Fibers, Tigran Baghdasaryan¹, Koen Vanmol¹, Hugo Thienpont¹, Francis Berghmans¹, Jurgen Van Erps¹; ¹*Vrije Universiteit Brussel, Belgium*. We used triangular cross-section multimode interference coupler, S-bends, and adiabatic tapers as building blocks to design and fabricate ultracompact 3D splitters for multi-core optical fiber with an insertion loss below -3 dB per channel.

IW3B.3 • 17:15

Highly Efficient TM Fundamental Mode Filter on InP Membrane, Dong Liang¹, Salim Abdi¹, Sander Reniers¹, Jos van der Tol¹, Kevin Williams¹, Yuqing Jiao¹; ¹*Eindhoven university of technology, Netherlands*. We experimentally demonstrate an efficient TM₀ filter on InP membrane. The filter exhibits an extinction ratio exceeding 34 dB for TM₀ and a loss lower than 0.5 dB for TE₀ at a wavelength of 1570 nm

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

IW3B.4 • 17:30

CubeSat Astrophotonics: Lower Cost, Space-Based Optical Astronomy Using Photonic Integrated Circuits, Tyler J. deLoughery^{1,2}, Clayton D. Lauzon^{1,2}, Kyle H. Sims^{1,2}, John Weber³, Wahab Almuhtadi^{2,1}, Ross Cheriton³; ¹*Carleton University, Canada*; ²*Faculty of Technology and Trades, Algonquin College, Canada*; ³*National Research Council Canada, Canada*. CubeSats are low-cost, space-based platforms for sensing instrumentation, where limits on size, weight and power limitations favour using photonic integrated circuit. We explore the design and performance of a proof-of-concept astrophotonic CubeSat for optical astronomy.

IW3B.5 • 17:45

High Dispersion in Hybridized Modes of Silicon Photonic Devices Compatible With a Foundry Platform, Archana Kaushalram¹, Jacob Hiesener¹, Clay A. Kaylor¹, Stephen Ralph¹; ¹*ECE, Georgia Institute of Technology, USA*. We report a peak dispersion of $\pm 10^5$ ps/nm/km with a bandwidth >35 nm in hybridized modes of a loaded-strip and loaded-slot waveguide with L-rails, structures that meet design rule checks of a commercial foundry.

IW3B.6 • 18:00 Invited

Image-to-Image Computer Vision for Advanced Nanophotonic Fabrication and Design, Dusan Gostimirovic¹, Odile Liboiron-Ladouceur¹; ¹*Department of Electrical and Computer Engineering, McGill University, Canada*. We present PreFab, an image-to-image computer vision system that predicts and corrects nanofabrication variations in complex integrated photonic circuits with fine features, enabling higher precision, improved yields, and accelerated development of next-generation photonic technologies.

16:30 -- 18:30

Room: 205B

SW3G • Radiative Cooling IV & Thermophotovoltaics & Perovskites (Joint SOLED/NOMA)

Presider: Canek Fuentes-Hernandez; Northeastern University, United States

SW3G.1 • 16:30 Invited

Nonlinear Optical Properties and Exciton Dynamics of Quantum Confined Halide Perovskites, Suchi Guha¹; ¹*University of Missouri-Columbia, USA*. Two-dimensional (2D) organic-inorganic halide perovskites emphasize a strong excitonic contribution, which enhances both linear and nonlinear optical properties. We present third harmonic generation and transient absorption studies from 2D and other quantum confined perovskites.

SW3G.2 • 17:00 Invited

Perovskite Single-Crystals: New Opportunities for Optoelectronics and Photonics, Rosanna Mastroia¹; ¹*CNR Nanotec, Italy*. Perovskite single-crystals emerged as a promising alternative to polycrystalline samples for optoelectronics. This talk highlights how the growth of perovskite single-crystals can be tailored to obtain specific features that fulfil the requirements of device integration.

SW3G.3 • 17:30 Invited

Bottom-Up Engineered Halide Perovskites for Optoelectronic Devices, Farnaz Niroui¹; ¹*Massachusetts Institute of Technology, USA*. This talk will introduce bottom-up engineering of designer materials with unique functionalities for emerging optoelectronic devices with a focus

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

on metal halide perovskites and their applications in nanoscale light-emitting devices and quantum light sources.

SW3G.4 • 18:00

Fabrication and Optimization of Highly Solar Reflective and Long-Wavelength Infrared (LWIR) Emissive Porous Polymers for Passive Daytime Radiative Cooling, Atousa Pirvaram¹, Paul G. O'Brien¹, Siu N. Leung¹; ¹*York University, Canada.*

This study explores the fabrication and optimization of micro- and nano-cellular poly(vinylidene fluoride-co-hexafluoropropylene) (PVDF-HFP) for passive daytime radiative cooling using phase inversion technique. By tailoring the water amount a high reflectance and a significant emittance are achieved, facilitating efficient cooling

SW3G.5 • 18:15 Postdeadline Submission

Optimal Thermophotovoltaics Using Bilayer Emitters, Paige Delsa¹, Mariama Rebello de Sousa Dias¹; ¹*University of Richmond, USA.* In thermophotovoltaics, components are required to function reliably under high temperatures. Here, we describe the optimal operating temperature, photovoltaic bandgap, and coating thickness of 283 bilayer emitters that are thermal stable beyond 1,800°C.

16:30 -- 18:30

Room: 206B

NpW3E • Special Symposium: The Future of Nonlinear Photonics

Presider: Ksenia Dolgaleva; University of Ottawa, Canada

NpW3E.1 • 16:30 Invited

An Embarrassment of Riches: What to Do With a Material 1 Million Times More Nonlinear Than Silica, Robert W. Boyd¹; ¹*University of Ottawa, Canada.*

Research reported in 2016 [Alam, De Leon, Boyd, *Science* 352, 795-797 (2016)] showed that materials such as indium tin oxide (ITO) display an unprecedentedly large nonlinear refractive index when excited at a frequency where the dielectric permittivity is nearly vanishing. The large nonlinear response of these epsilon-near-zero (ENZ) materials, some million times larger than that of silica glass, suggest that these materials would prove extremely useful in the development of new classes of photonic devices. In this contribution, we review the history of the understanding of nature of the nonlinear response of ENZ materials and the development of ENZ-based photonic devices. Specific investigations over the past eight years have led to increased understanding of topics including (1) the origin of the enhancement of the nonlinear optical response, (2) the possibility of using nanofabrication methods to create ENZ behavior at any specified wavelength, and (3) the exploration of what interactions and devices are enabled through use of ENZ materials. In this contribution, we review the recent history of the development of the field of ENZ science and of the development of ENZ applications.

NpW3E.2 • 16:50 Invited

Progress Towards Silicon Nitride as a $\chi^{(3)}$ and $\chi^{(2)}$ Nonlinear Optics Platform, Jaime Cardenas¹; ¹*University of Rochester, USA.* Silicon nitride is the leading on-chip platform for $\chi^{(3)}$ nonlinear optics. It has recently emerged as candidate for $\chi^{(2)}$ nonlinear processes through optical and electrical poling. We review recent progress in silicon nitride $\chi^{(2)}$ photonics.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

NpW3E.3 • 17:10 Invited

Title to be Announced, Alexander L. Gaeta¹; ¹*Columbia University, USA*. Abstract not available.

NpW3E.4 • 17:30 Invited

Extremely Non-Degenerate Nonlinear Optics: Dedicated to the Memory of Mansoor Sheik-Bahae, Eric W. Van Stryland¹; ¹*University of Central Florida, CREOL, USA*. Mansoor et.al. introduced Z-scan 35 years ago (10^4 citations) which led to understanding Kramers-Kronig for bound-electronic nonlinearities, which led to establishing and understanding the large enhancement of nonlinearities for Extremely Non-Degenerate (END) nonlinear optics.

NpW3E.5 • 17:50 Invited

Perspectives on the Future of Nonlinear Photonics: the Promise of Well-Established and Novel Materials, Nathalie Vermeulen¹; ¹*Vrije Universiteit Brussel, Belgium*. The rise of low-dimensional materials, meta-materials, etc. has expedited nonlinear-optics research. Meanwhile, fabrication technologies for well-established materials have significantly advanced. This talk will explore the implications of both developments for the future of nonlinear photonics.

16:30 -- 18:30

Room: 205C

NoW3D • Laser Assisted Processing of Optics

Presider: Francois Chenard; IRflex Corporation, United States

NoW3D.1 • 16:30 Invited

Laser-Assisted Deposition of Oxide Layers for Transparency in Different Wavelength Ranges, Yahya Bougdid¹, Gunjan Kulkarni¹, Francois Chenard², Chandrika J. Sugrim⁴, Ranganathan Kumar³, Aravinda Kar¹; ¹*University of Central Florida, CREOL, USA*; ²*IRflex Corporation, USA*; ³*University of Central Florida, USA*; ⁴*Naval Air Warfare Center, Aircraft Division, USA*. CO₂ laser-assisted deposition of transparent chalcogenide As₂S₃ glass and TiO₂ films is reported. A theoretical model is developed to select the laser sintering parameters. The optical properties of transparent As₂S₃ and TiO₂ coatings are investigated.

NoW3D.2 • 17:00 Invited

Digital Glass Forming of Optics and Photonics, Edward Kinzel¹; ¹*University of Notre Dame, USA*. CO₂ lasers are used for digital glass forming. This produces surface heating that slowly diffuses through the filament. We investigate using a NIR laser and glass doping for faster printing of optics and photonics.

NoW3D.3 • 17:30 Invited

Ultrafast Lasers Enabled Optical Figuring and 3D Writing, Jie Qiao¹; ¹*Rochester Institute of Technology, USA*. Abstract not available.

NoW3D.4

Withdrawn

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

16:30 -- 18:30

Room: 2104

NoW3H • Optical Sensors, Biosensors, and Imaging

Presider: Alon Gorodetsky; University of California Irvine, United States

NoW3H.1 • 16:30 Invited

Title to be Announced, Tian Li¹; ¹*Purdue University, USA*. Abstract not available.

NoW3H.2 • 17:00

A Multiplexed Quantitative Spectroscopy Method to Measure Spectrally Overlapping Metabolic Fluorophores, Victoria W. D'Agostino¹, Michelle Kwan¹, Megan Madonna¹, Brian Crouch¹, Nimmi Ramanujam¹; ¹*Duke University, USA*. We have developed and validated an *in vivo* quantitative spectroscopy method to measure spectrally overlapping fluorophores relevant to cancer metabolism using an inverse Monte Carlo algorithm and linear spectral unmixing.

NoW3H.3 • 17:15

Terahertz Time-Domain Derivative Spectroscopy of Fructose Using a MEMS Piezo Speaker, Behnoosh Meskoob¹, Mathieu Gratuze¹, Gabriel Gandubert¹, Xavier Ropagnol¹, Frederic Nabki¹, Francois Blanchard¹; ¹*École de Technologie Supérieure, Canada*. This paper presents the implementation of MEMS to convert Terahertz Time-Domain Spectroscopy (THz-TDS) to the Time-Domain Derivative Spectroscopy (THz-TDDS). This THz signal modulation is achieved without losing the spectroscopy ability for material characterization.

NoW3H.4 • 17:30

Feature-Dependent Accuracy in Classification Models for Ultrafast THz Spectroscopy Using Frequency Selective Surfaces, Rejeena R Sebastian¹, Redwan Ahmad¹, Jonathan Lafrenière-Greig¹, Xavier Ropagnol^{1,2}, Francois Blanchard¹; ¹*École de technologie supérieure ÉTS, Canada*; ²*Institut National de la recherche scientifique (INRS-EMT), Canada*. This work analyzes data from a novel setup for ultrafast selective multispectral Terahertz (THz) spectroscopy using Frequency Selective Surface (FSS). A comparative evaluation of the performance of four different classification models is conducted and a feature-accuracy mapping is done for optimization.

NoW3H.5 • 17:45

Spiropyran-Doped Poly(Dimethylsiloxane) Optical Waveguides for UV Sensing, Camila Zimmermann¹, Koffi N. Amouzou¹, Dipankar Sengupta¹, Aashutosh Kumar¹, Nicole Demarquette¹, Bora Ung¹; ¹*École de technologie supérieure ÉTS, Canada*. Novel spiropyran-doped PDMS optical waveguides were fabricated. Their UV sensing response was investigated by monitoring changes in transmitted optical power when coupled to a 633-nm HeNe laser.

NoW3H.6 • 18:00

Novel Porous-Cladding Polydimethylsiloxane Optical Waveguide for Biomedical Pressure Sensing Applications, Koffi N. Amouzou¹, Camila Zimmermann¹, Bora Ung¹, Dipankar Sengupta¹, Normand Gravel¹, Jean-Marc Lina¹, Bora Ung¹; ¹*École de technologie supérieure, Canada*. We report a new concept of pressure sensor made from polydimethylsiloxane solid core and porous cladding that operates through frustrated total internal reflection. A high sensitivity to transverse compression of 0.22%/dB optical losses is demonstrated.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

16:30 -- 18:30

Room: 2101

NeW3C • Coherent Access Networks

Presider: Domaniç Lavery; Infinera Corporation, United Kingdom

NeW3C.1 • 16:30 Invited

Coherent PON for Optical Access Networks Evolution Beyond 50G-PON, Giuseppe Talli¹; ¹*Huawei, Germany*. Coherent PON is a promising candidate for the next generation VHSP-PON systems, which could support net data rates beyond 200Gb/s per wavelength, maintain coexistence with previous generations, and enable new services.

NeW3C.2 • 17:00 Invited

Coherent Optical Technologies Shaping the Evolution of Passive Optical Networks, Zhensheng Jia¹; ¹*CableLabs, USA*. This paper discusses the evolution of PON technologies by ITU-T and IEEE. It evaluates the progress and limitations of IM-DD PONs, and presents the drivers for longer reach and higher split coherent PONs. The paper also explores key technology developments and options to simplify coherent designs for cost reduction.

NeW3C.3 • 17:30

Supporting 100 Gb/s and 200Gb/s Optical Access for PON Using Coherent Technologies, Mark E. Laubach¹; ¹*Ciena Corporation, USA*. Demand for higher speed subscriber networks coupled with developing ecosystems for coherent point-to-point technologies suggest cost-effective re-use for 100 Gb/s and 200 Gb/s for the standardization of Coherent PON.

16:30 -- 18:30

Room: 207

SoW3F • Active Fibers, Lasers & Amplifiers

Presider: Martin Rochette; McGill University, Canada

SoW3F.1 • 16:30 Invited

Withdrawn

SoW3F.2 • 17:00

All Fiber Mid-Infrared Ring Cavity Laser, Nasrollah Karampour¹, Gebrehiwot T. Zeweldi¹, Md. Hosne Mobarok Shamim¹, Martin Rochette¹; ¹*McGill university, Canada*. We demonstrate the first all-fluoride mid-infrared ring cavity laser, comprising a single-mode ZBLAN optical fiber coupler and an Er: ZBLAN gain fiber. The laser exhibits continuous-wave emission at a wavelength of 2.7-2.8 μm .

SoW3F.3 • 17:15

Mode-Locking Using Polarization Dependent Fluoride Optical Fiber Coupler, Gebrehiwot T. Zeweldi¹, Martin Rochette¹; ¹*McGill University, Canada*. We make the first demonstration of a polarization-dependent single-mode fluoride-based optical fiber coupler. The practicality of this mid-infrared compatible coupler is shown with the successful realization of an all-fiber mode-locked ring cavity laser.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

SoW3F.4 • 17:30

Improved Emission Lifetime via Thermal Annealing for Ytterbium-Doped High-Power Fiber Lasers, Siyuan Wang¹, Bailey Meehan², Thomas Wade Hawkins², John Ballato², Peter D. Dragic¹; ¹*University of Illinois Urbana Champaign, USA*; ²*Clemson University, USA*. Through a series of thermal annealing experiments, it is shown that Yb³⁺ emission properties could be further optimized, particularly as relates to the quantum efficiency. Achieving near-unity quantum efficiency is paramount to continued power scaling.

SoW3F.5 • 17:45

Effects of Package Layout on Active Fibre Absorption, Mihai-Stefan Merlas¹, Natasha Vukovic¹, Christophe A. Codemard², Michalis N. Zervas¹; ¹*University of Southampton, United Kingdom*; ²*TRUMPF Lasers UK, United Kingdom*. We investigate numerically the impact of layout in the absorption in double-clad fibres. We find that the local curvature and its variation along the fibre length are the key parameters, defining the total pump absorption.

SoW3F.6 • 18:00

Larger Cladding Diameter for Improved Power Conversion Efficiency in Cladding Pumped Super L-Band Amplifiers, Saber Jalilpiran¹, Hamed Rabbani¹, Jacques Lefebvre¹, Leslie Rusch¹, Younès Messaddeq¹, Sophie LaRochelle¹; ¹*Centre d'optique, photonique et laser (COPL), Laval University, Canada*. We compare pump power requirements of two fibers having respective cladding diameters of 125 mm and 200 mm in cladding-pumped amplifiers. The Er³⁺:Yb³⁺ co-doped alumino-phospho-silicate fibers were optimized for super L-band operation.

SoW3F.7 • 18:15

Impact of Annealing and Fictive Temperature on Brillouin and Raman Scattering Spectra in Yb-Doped Fibers, Siyuan Wang¹, Bailey Meehan², Thomas Wade Hawkins², John Ballato², Peter D. Dragic¹; ¹*University of Illinois Urbana Champaign, USA*; ²*Clemson University, USA*. The impacts of thermal annealing on Brillouin and Raman scattering in Yb-doped fibers are studied and explained by mechanical stress relief, ion diffusion, and glass relaxation. The results facilitate improved performance for high-power laser applications.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Thursday, 1 August

08:00 -- 10:00

Room: 206A

BTh1A • Applications of FBG and Laser Written Devices

Presider: Christophe Caucheteur; Universite de Mons, Belgium

BTh1A.1 • 08:00 Invited

Ultrashort Pulse Written VBGs: Achievements and Next Step, Daniel Richter¹, Malte P. Siems¹, Ria G. Krämer¹, Georg R. Schwartz¹, Abdolnaser Ghazagh¹, Stefan Nolte^{1,2}; ¹*Institute of Applied Physics, Abbe Center of Photonics, Friedrich Schiller University Jena, Germany*; ²*Fraunhofer Institute for Applied Optics and Precision Engineering, Center of Excellence in Photonics, Germany*. With ultrashort pulsed VBGs, the already impressive application areas of commercial VBGs can be even further expanded by overcoming process-related material limitations. Achievements, performance, current challenges and future applications will be highlighted.

BTh1A.2 • 08:30

Dual Mode-Comb Plasmonic Optical Fiber Sensing, Gabriel E. Villatoro-Perez², Médéric Loyez³, Joel Villatoro⁴, Christophe Caucheteur⁵, Jacques Albert¹; ¹*Electronics, Carleton University, Canada*; ²*Optics, National Institute of Astrophysics Optics and Electronics, Mexico*; ³*Proteomics and Microbiology, University of Mons, Belgium*; ⁴*Ingeniería de Comunicaciones UPV/EHU, University of the Basque Country UPV/EHU, Spain*; ⁵*Electromagnetism and Telecom, University of Mons, Belgium*. A partially gold-coated tilted FBG is proposed for self-referenced plasmonic sensing. The device exhibits two interleaved combs of resonances with unpolarized light; one comb is used as a reference and the other as a sensor.

BTh1A.3 • 08:45

Distributed Magnetic and Current Sensing Through Enhanced-Signal Random Bragg Grating, Antoine Leymonerie¹, Olivier Bélanger¹, Jean-Sébastien Boisvert¹, Sébastien Loranger¹; ¹*Electrical Engineering, Polytechnique Montréal, Canada*. We demonstrate the advantage of random fiber Bragg grating for distributed magnetic and current sensing. The 40 dB signal increase of the gratings allows a measurement down to 10 mT with 8 cm spatial resolution.

BTh1A.4 • 09:00

Multi-Channel and Dual-Range Spectrum Analyzer for Low-Cost Parallel TFBG Sensing, Julian Nicolai¹, Hubert Jean-Ruel¹; ¹*Department of Electronics, Carleton University, Canada*. An inexpensive spectrometer scheme for multiplexed TFBG biosensors is proposed. Multiple channels and two spectral windows are combined on a CMOS camera to measure with high-resolution a selection of cladding modes and the Bragg peak.

BTh1A.5 • 09:15

Fiber Bragg Grating Based Hydrogen Leak Detection, Korina Hartmann¹, Remco Nieuwland¹; ¹*United Fiber Sensing B.V., Netherlands*. A Fiber Bragg grating (FBG)-based hydrogen sensor coated with a catalytic layer was developed, detecting H₂ from 0.1 vol.% with response times of 3s. Experimental verification showed safe usage above the lower explosion limit (LEL).

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

BTh1A.6 • 09:30 Invited

The Road to a Practicable Magnetless Integrated Optical Isolator, Jerome Lapointe¹, Cedrik Coia², Albert Dupont¹, Réal Vallée¹; ¹*Center for Optics, Photonics and Lasers, Canada*; ²*Aeponyx, Canada*. Since the 1990s, optical isolators have hindered full integration of photonic chips. Here, we describe challenges and a successful solution: an fs-laser-inscribed low-loss magnetless broadband integrated isolator. Additionally, we discuss potential routes to mass production.

08:00 -- 10:00

Room: 205A

ITh1B • Novel Integrated Materials I

Presider: Christian Haffner; Interuniversity Microelectronics Center, Belgium

ITh1B.1 • 08:00 Tutorial

Advanced Integrated Photonics on Lithium Niobate, Sasan Fathpour¹; ¹*University of Central Florida, CREOL, USA*. Abstract not available.

ITh1B.2 • 09:00

Four-Wave Mixing in Highly Nonlinear Silicon Slot Waveguides With a Crystal Violet Cladding, Devika P. Nair¹, Michael Menard¹; ¹*École de technologie supérieure, Canada*. We introduce a silicon slot waveguide encased in a highly nonlinear cladding containing crystal violet molecules. We measured the four-wave mixing efficiency and results show potential for implementing compact high-performance nonlinear optical devices.

ITh1B.3 • 09:15

The Effect of a Protective ALD Layer on the Stability of Electro-Optic Polymer, Marwan Albarghouti¹, Mateo Powell Serrano¹, Jasper Drisko¹, Patrick Riedel¹, Gannon Kehe¹, Ginnelle Ramann¹, Xiaoyue P. Huang¹, Brenden Basica¹, John Zyskind¹, Zhiming Liu¹; ¹*Lightwave Logic, Inc., USA*. We describe a method to improve the photo and thermal stability of electro-optic polymers. Our results show that degradation is reduced by encapsulating the EOP with a layer that is impermeable to molecular oxygen O₂

ITh1B.4 • 09:30 Invited

Near-Zero-Power Consumption Chalcogenide Phase Change Materials Photonic Devices for OFPGA Applications, Hongtao Lin¹; ¹*Zhejiang University, China*. Abstract not available.

08:00 -- 10:00

Room: 206B

NpTh1E • Frequency Combs and Spectral Broadening

Presider: Sile Nic Chormaic; Okinawa Inst of Science & Technology, Japan

NpTh1E.1 • 08:00 Invited

Title to be Announced, Zaijun Chen¹; ¹*University of Southern California, USA*. Abstract not available.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

NpTh1E.2 • 08:30

Coherent Generation of Ultra-Stable Smart Frequency Combs, Celine Mazoukh¹, Luigi Di Lauro¹, Imtiaz Alamgir¹, Bennet Fischer^{1,2}, Nicolas Perron¹, A Aadhi³, Armaghan Eshaghi⁴, Brent Little⁵, Sai Chu⁶, David J. Moss⁷, Roberto Morandotti¹; ¹*INRS-EMT, Canada*; ²*Leibniz Institute of Photonic Technology, Germany*; ³*Centre for Nanophotonics, Queen's University, Canada*; ⁴*Huawei Technologies Canada, Canada*; ⁵*QXP Technologies, China*; ⁶*City University of Hong Kong, Hong Kong*; ⁷*Optical Sciences Centre, Swinburne University of Technology, Australia*. We present a novel smart method to customize microcomb state generation in microring resonators pumped with a continuous-wave laser, using genetic algorithms to identify optimal experimental parameters for coherent state generation.

NpTh1E.3 • 08:45

Control of Frequency Comb Spacing via Self-Crystallization of Dark Vectorial Solitons in Kerr Resonators, Graeme N. Campbell^{1,2}, Lewis J. Hill², Pascal Del'Haye^{2,3}, Gian-Luca Oppo¹; ¹*Department of Physics, University of Strathclyde, United Kingdom*; ²*Max Planck Institute for the Science of Light, Germany*; ³*Department of Physics, Friedrich-Alexander-Universitat Erlangen-Nurnberg, Germany*. Dark vectorial solitons in Kerr resonators with normal dispersion can spontaneously self-organize into regularly spaced crystals resulting in greater power and spacing of the frequency comb lines with increasing soliton numbers.

NpTh1E.4 • 09:00

Frequency-Offset Kerr Soliton Comb Generation in a Dispersion-Shifted Fiber Fabry-Perot Resonator, Yiqing Xu^{1,2}, Matthew Macnaughtan^{1,2}, Zongda Li^{1,2}, Xiaoming Wei³, Zhongmin Yang³, Stephane Coen^{1,2}, Miro Erkintalo^{1,2}, Stuart Murdoch^{1,2}; ¹*The Dodd-Walls Centre for Photonic and Quantum Technologies, New Zealand*; ²*Department of Physics, University of Auckland, New Zealand*; ³*School of Physics and Optoelectronics, South China University of Technology, China*. We present a study of frequency-offset soliton combs that arise when a Kerr resonator is driven by a desynchronized pulsed field. We experimentally observe these offset-combs in a Fabry-Perot resonator constructed from dispersion-shifted optical fiber.

NpTh1E.5 • 09:15

Tunable Timescale Mode-Locked Laser Based on a Nested Microring Resonator, Imtiaz Alamgir¹, A Aadhi¹, Luigi Di Lauro¹, Pavel Dmitriev¹, Nicolas Perron¹, Celine Mazoukh¹, Bennet Fischer², Piotr Roztock^{1,3}, Cristina Rimoldi^{1,4}, Mario Chemnitz^{1,2}, Armaghan Eshaghi⁵, Evgeny Viktorov⁶, Anton Kovalev⁶, Brent Little⁷, Sai Chu⁸, David J. Moss⁹, Roberto Morandotti¹; ¹*Energie Matériaux Télécommunications, Institut national de la recherche scientifique, Canada*; ²*Leibniz Institute of Photonic Technology, Germany*; ³*Ki3 Photonics Technologies, Canada*; ⁴*Dipartimento di Elettronica e Telecomunicazioni, Politecnico di Torino, Italy*; ⁵*Huawei Technologies Canada, Canada*; ⁶*ITMO University, Russian Federation*; ⁷*QXP Technology Inc., China*; ⁸*City University of Hong Kong, China*; ⁹*Swinburne University of Technology, Australia*. We demonstrate an active mode-locked laser based on a nested microring resonator configuration that generates and switches between different repetition rates. This technique offers a versatile solution for spectroscopy, micromachining, and telecommunications applications.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

08:00 -- 10:00

Room: 205B

NoTh1G • Chalcogenides and Nonlinear Materials

Presider: Brandon Shaw; Naval Research Laboratory, United States

NoTh1G.1 • 08:00 Invited

On the Temperature Dependence of Photo-Structural Changes in Chalcogenide Glasses, Pierre Lucas¹; ¹*University of Arizona, USA*. This presentation will show that while photo-structural changes in chalcogenide glasses are unquestionably optically induced, there is also clear evidence that many of these changes are thermally facilitated even at moderate laser power.

NoTh1G.2 • 08:30

Demonstration of Optical Refrigeration in SiO₂ – Al₂O₃ – Lu₂O₃ Yb-Doped Glasses, Thomas Meyneng^{1,2}, Jyothis Thomas¹, Nicolas Grégoire², Morency Steeve², Philippe Labranche², Jean-Sebastien Boisvert¹, Younès Messaddeq², Raman Kashyap¹; ¹*Department of electrical engineering, Polytechnique Montréal, Canada*; ²*COPL, Université Laval, Canada*. This work demonstrates the applications of Lu₂O₃-Al₂O₃-SiO₂ Yb-doped glasses for optical refrigeration. Samples prepared by modified chemical vapor deposition, combined with solution-doping, exhibits near-unity quantum efficiency, with Yb-content ranging from 10²⁵ to 10²⁶ m⁻³.

NoTh1G.3 • 08:45

Tunable Continuous Wave Tm³⁺-Doped Gallium-Rich BGG Glass Laser, Stanislav Leonov¹, Théo Guérineau¹, Martin Bernier¹, Younès Messaddeq¹, Réal Vallée¹; ¹*Center for Optics, Photonics and Lasers (COPL), Université Laval, Canada*. We present a room-temperature tunable continuous wave laser based on Tm³⁺-doped Ga-rich BGG glass. The laser wavelength was continuously tuned from 1855 to 1965 nm, representing a tuning range of 110 nm.

NoTh1G.4 • 09:00

Comparison of Laser Induced Damage Thresholds of CdSiP₂ and BaGa₂GeSe₆ at 1064 nm for Different Spot Sizes, Shekhar Guha¹, Kevin Cissner¹, Alexander Carson¹, Kevin Zawilski², Valentin Petrov³; ¹*US Air Force Research Laboratory, USA*; ²*BAE Inc., USA*; ³*Max Born Institut, Germany*. Damage threshold values of CdSiP₂ and BaGa₂GeSe₆ under 5 ns duration 1064 nm laser exposure were measured for various spot sizes.

NoTh1G.5 • 09:15

Tuning Effective Optical Nonlinearities of Overlooked Glass-Forming Ionic Liquid Crystals, Valentyn Rudenko², Anatolii Tolochko², Svitlana Bugaychuk², Dmytro Zhulai², Gertruda Klimusheva², Galina Yaremchuk³, Tatyana Mirnaya³, Yuriy Garbovskiy¹; ¹*Central Connecticut State University, USA*; ²*Institute of Physics of National Academy of Sciences of Ukraine, Ukraine*; ³*V.I. Vernadsky Institute of General and Inorganic Chemistry of National Academy of Sciences of Ukraine, Ukraine*. This paper reports the modification of nonlinear-optical properties of very common yet often overlook glass-forming mesogenic materials (metal alkanoates) by using several types of nanoparticles including metal (silver and gold), bimetallic, and carbon dots.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

NoTh1G.6 • 09:30 Invited

Mid Infrared Photonics on Thin-Film Lithium Niobate, Mengjie Yu¹; ¹*University of Southern California, USA*. Abstract not available.

08:00 -- 10:00

Room: 205C

NoTh1D • Nanophotonics

Presider: Jonathan Hu; Baylor University, United States

NoTh1D.1 • 08:00 Invited

Optical Positioning and Linking Using Optical Tweezers for 3D Nanofabrication, Euan McLeod¹; ¹*University of Arizona, USA*. Optical tweezers can accurately position nanoparticles of a wide range of materials in complex geometries. Assembly is automated using computer control. Applications include the fabrication of grating arrays for free-space coupling into microtoroidal optical sensors.

NoTh1D.2 • 08:30 Invited

Nanowire Terahertz Detectors: From Laboratory to Industrial Scale, Kun Peng¹, Nicholas Morgan², Ford Wagner¹, Thomas Siday¹, Chelsea Xia¹, Didem Dede², Victor Boureau³, Valerio Piazza², Anna Fontcuberta i Morral^{2,4}, Michael Johnston¹; ¹*Department of Physics, University of Oxford, United Kingdom*; ²*Institute of Materials, EPFL, Switzerland*; ³*Interdisciplinary Centre for Electron Microscopy, EPFL, Switzerland*; ⁴*Institute of Physics, EPFL, Switzerland*. Nanowires are advantageous for the phase-sensitive coherent detection of terahertz radiation by enabling the measurement of the full terahertz polarization state. However, the development of parallel-nanowire detectors or large-scale imaging arrays is challenging. Our work delivered solutions to this challenge.

NoTh1D.3 • 09:00 Invited

UV to Mid-IR Single Photon Detection Using Superconducting Fractal Nanowires, Vidur Raj¹, Fiheon Imroze¹, Dmitry Morozov¹, Ciaran T. Lennon¹, Gregor G. Taylor², Martin Weides¹, Robert H. Hadfield¹; ¹*University of Glasgow, United Kingdom*; ²*EPFL, Switzerland*.

Superconducting nanowire single photon detectors (SNSPDs) have emerged as a preferred single photon detection technology for photonic quantum technology applications. Here, we show fractal nanowire SNSPDs operating from up to mid-IR regime, with saturating internal quantum efficiency.

NoTh1D.4 • 09:30

Withdrawn

NoTh1D.5 • 09:45

Tunable Nanophotonic Materials for Multispectral Reconfigurability, Yujie Luo¹, Thomas Christensen², Ognjen Ilic¹; ¹*University of Minnesota Twin Cities, USA*; ²*Technical University of Denmark, Denmark*. We show that active metasurface networks can accurately emulate and switch between complex spectral profiles, such as those of gases. This multispectral reconfigurability applies to coupled networks in active materials, including 2D or phase-change materials.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

08:00 -- 10:00

Room: 2101

NeTh1C • Wireless, Transport, and Sensing in Networks

Presider: Xiaodan Pang, KTH Royal Institute of Technology, Sweden

NeTh1C.1 • 08:00 Invited

Withdrawn

NeTh1C.2 • 08:30 Invited

Enabling Sub-THz 6G Wireless Systems Using Photonic Technologies, Liam P. Barry¹, Amol Delmade¹, Simon Nellen², David Coffey³, Cristian Vargas², Alison Kearney^{3,1}, Robert Kohlhaas², Martin Schell², Frank Smyth³; ¹*Dublin City University, Ireland*; ²*Fraunhofer Institute for Telecommunications, Heinrich Hertz Institute (HHI), Germany*; ³*Pilot Photonics, Ireland*. The key elements of an optical heterodyne analog radio-over-fiber link, i.e. optical source, detector technology and impairment compensation techniques, are analyzed for high frequency (180-260 GHz) sub-THz signal generation in 6G wireless systems

NeTh1C.3 • 09:00 Invited

Hybrid Digital PAM and 60 GHz Analog Radio Over Fiber Optical Spectrum as a Service Applications, Devika Dass¹; ¹*University of Dublin Trinity College, Ireland*. We experimentally study transmission of digitally multiplexed 15 Gb/s PAM-8, wideband 16-QAM OFDM (WiGig) and narrowband 64-QAM OFDM (5G NR) waveforms over a 10 km fiber, undergoing up-conversion to 60 GHz and achieving FEC limited performance.

NeTh1C.4 • 09:30

Optical Frequency Comb Generation for DWDM Free Space Optical Communication Links, Mahrokh Avazpour¹, Narmada Rajaram², Ahmad Atieh^{2,3}, Liam P. Barry¹; ¹*Dublin City University, Ireland*; ²*Department of Electrical and Computer Engineering, University of Ottawa, Canada*; ³*Optiwave System Inc., Canada*. This paper presents DWDM-based FSO system with 42 channels generated via SOA-Fiber Loop Modulation. System performance is assessed using BER and eye diagrams considering bit rate, and FSO range applicable for high-capacity FSO link.

08:00 -- 10:00

Room: 2104

SpTh1H • Coherent Technologies II

Presider: Dora van Veen; Nokia Corporation, United States

SpTh1H.1 • 08:00 Invited

Programmable Coherent Linear Processor for Machine Learning and Optical Communications, Mitsumasa Nakajima¹, Kohei Ikeda^{2,3}, Kohki Shibahara⁴, Akira Kawai⁴, Masaya Notomi^{2,3}, Takayuki Kobayashi⁴, Yutaka Miyamoto⁴, Toshikazu Hashimoto¹; ¹*NTT Device Technology Labs., Japan*; ²*NTT Nanophotonic Center, Japan*; ³*NTT Basic Research Labs., Japan*; ⁴*NTT Network Innovation Labs., Japan*. We demonstrate linear algebra processors on integrated photonic circuits. Applications to a machine learning accelerator and pre-processor for optical communications are demonstrated by implementing recurrent, convolutional, and unitary processors on photonic chips.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

SpTh1H.2 • 08:30

Add-Drop Multiplexing for Full Spectrum WDM NFDW Transmission Systems Using Spectral Overlap, Olaf Schulz¹, Alvaro Moscoso-Mártir², Jeremy Witzens², Stephan Pachnicke¹; ¹*Kiel University, Germany*; ²*RWTH Aachen University, Germany*. We present add-drop multiplexing for full spectrum modulated WDM nonlinear frequency division multiplexed transmission systems, in which spectral overlap between channels is used to eliminate nonlinear inter-channel crosstalk.

SpTh1H.3 • 08:45

From Analog Coherent Optics to Linear Drive Pluggable Optics: Lessons Learnt, Naim Ben-Hamida¹, Shahab Oveis Gharan¹, Ahmad Abdo¹, Bilal Riaz¹; ¹*Ciena Canada Inc., Canada*. Linear drive pluggable optics (LPO) forms a paradigm shift in the short-reach optical switching space. In here, we go through the trajectory from the CFP2-Analog Coherent Optics (ACO) to LPO and discuss the challenges to tackle as data rates increase.

SpTh1H.4 • 09:00

Interlocking of Mode-Locked Lasers Utilizing Optical Communication Equipment, Ido Attia^{1,2}, Dan Sadot^{1,2}; ¹*Ben-Gurion University of the Negev, Israel*; ²*CyberRidge, Israel*. Interlocking of mode-locked lasers (MLLs) has several applications in the optical communication field and beyond. This work uses an on-chip, integrated coherent receiver (ICR) for successfully interlocking two 80GHz MLLs without exploiting the ICR high-speed outputs.

SpTh1H.5 • 09:15 Invited

Integrated Comb Lasers for Coherent Transceiver Scaling, Frank Smyth¹; ¹*Pilot Photonics, Ireland*. Abstract not available.

SpTh1H.6 • 09:45

Feedforward Phase Noise Compensation of Free-Running DFB Lasers Heterodyned for sub-THz Generation Without Optical Filtering, Kyungmin Woo¹, Hoon Kim¹; ¹*KAIST, Korea (the Republic of)*. We experimentally demonstrate the feedforward compensation of phase fluctuations of two free-running DFB lasers heterodyned for 166-GHz signal without any optical filtering. We achieve the SSB phase noise of -80 dBc/Hz at 100-kHz offset frequency.

08:00 -- 10:00

Room: 207

SoTh1F • Fibers and Devices for Biomedical and Sensing Applications I

Presider: Natalie Wheeler; University of Southampton, United Kingdom

SoTh1F.1 • 08:00 Invited

Novel Fiber Optic Components for Biomedical Imaging and Sensing, Caroline Boudoux¹; ¹*Polytechnique Montréal, Canada*. Abstract not available.

SoTh1F.2 • 08:30

Distributed Hydrogen Sensing and Leak Detection Using Draw-Tower Fabricated Optical Fiber, sandy alomari¹, Kenny Hey Tow¹, Joao Pereira¹, Ari Antikainen¹, Tedros Weldehawariat¹, Korina Hartmann², Remco Nieuwland², Åsa Claesson¹; ¹*RISE Fiberlab, RISE Research Institutes of Sweden, Sweden*; ²*United Fiber Sensing, Netherlands*. A long length distributed

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

hydrogen sensor was demonstrated for the first time, using a draw-tower fabricated optical fiber with a sensor coating that reacts exothermically in the presence of hydrogen in air.

SoTh1F.3 • 09:00

Anti-Resonant, Hollow-Core Fiber Enhanced Raman Gas Spectroscopy With a 520nm

Edge-Emitting Laser Diode, Ian A. Davidson¹, Thomas Kelly¹, Peter Horak¹, David J. Richardson¹, Francesco Poletti¹, Natalie V. Wheeler¹; ¹*University of Southampton, United Kingdom*. Hollow-core fiber enhanced Raman spectroscopy is a highly versatile technique that normally utilizes narrow linewidth pump-lasers. Here a low-cost, ~1nm linewidth, edge-emitting laser diode is used resulting in a potentially more compact and energy-efficient system.

SoTh1F.4 • 09:15

Tapering Space-Division Multiplexing Fibers for Multi-Parameter Sensing

Liudmila Silanteva¹, Vincent V. Vliet¹, Menno V. Hout¹, Chigo Okonkwo¹, Thomas Bradley¹; ¹*Eindhoven University of Technology, Netherlands*. We report on simulations of coupled-core fibers tapered to a diameter $\approx 1.2 \mu\text{m}$, resulting in mode field diameters of $\approx 1.3 \mu\text{m}$ (LP01) and $\approx 2 \mu\text{m}$ (LP11) respectively for enhancing the field for a multi-parameter sensor.

SoTh1F.5 • 09:30

Optical Fiber SERS Sensors for Salmonella Detection in Poultry Products

Mai Abuhelwa¹, Arshdeep Singh¹, Jiayu Liu¹, Amit Morey², Lakshmi Kantha Channaiah¹, Mahmoud Almasri¹; ¹*University of Missouri-Columbia, USA*; ²*Auburn University, USA*. This paper introduces SERS sensors with nanoantenna arrays on an optical fiber and 3D printed microstructure, detecting 1-3 cells/ml in 10 minutes. It enhances signal-to-noise ratio through increased surface area, unpublished.

SoTh1F.6 • 09:45

THz Negative Curvature Fiber Sensor Design for Blood Constituent Analysis

Julia Ward¹, Ethan Neidt¹, Riley Como¹, Ahmet E. Akosman¹; ¹*Roger Williams University, USA*. A unique hollow-core negative curvature fiber design to achieve broadband sensing of blood constituents in the THz range is proposed. Numerical investigations indicate operational bandwidths exceeding 0.6 THz and simultaneous selectivity for different constituents.

10:30 -- 12:30

Room: 206A

BTh2A • Optical Fiber Sensing

Presider: Patrice Mégret; Universite de Mons, Belgium

BTh2A.1 • 10:30 Invited

Optical Fiber Sensors for Volatile Organic Compound Vapors

Eleni Grantzioti¹, Panagiotis Kleitsiotis^{1,2}, Emmanouil Gagaoudakis¹, Vassilios Binas^{1,3}, Stavros Pissadakis¹, Maria Konstantaki¹; ¹*FORTH-IESL, Greece*; ²*Physics Department, University of Crete, Greece*; ³*Chemistry Department, Aristotle University of Thessaloniki, Greece*. A review of optical fiber sensors utilizing tilted Bragg gratings, long-period gratings, or Fabry-Perot interferometer coupled with metal oxide or polymeric sensing overlays/cavities for the detection of methanol, ethanol, isopropanol and acetone vapors.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

BTh2A.2 • 11:00

Developing Robust Optical Fibre Sensors for use in Hostile Sewer Environments, Lachlan Anderson¹, Peter Dekker¹, Heriberto Bustamante², Thomas Kuen³, Michael J. Withford¹, Martin Ams¹; ¹*Macquarie University, Australia*; ²*Sydney Water, Australia*; ³*Melbourne Water, Australia*. We report robust fibre Bragg grating (FBG) sensors that optically measure environmental conditions in concrete wastewater networks over long periods. We also demonstrate an optical fibre dew point sensor.

BTh2A.3 • 11:15

Spectral Detangling of Plasmonic Fiber Bragg Grating Biosensors, Hadrien Fasseaux¹, Christophe Caucheteur¹, Médéric Loyez¹; ¹*Universite de Mons, Belgium*. Plasmonic tilted fiber Bragg gratings are highly sensitive refractometers and biosensors. The detection occurs through the detangling of intricate comb-like spectral signatures and various demodulation techniques have been proposed. We study here their relative performance.

BTh2A.4 • 11:30

Optical Vernier Effect Based on High Order Fiber Bragg Grating, Romain Cotillard¹, Nicolas Roussel¹; ¹*CEA, France*. The use of the Optical Vernier effect to increase sensitivity to a physical parameter is well known in the field of Fabry-Perot cavities. We propose here a simplified application with fiber Bragg gratings.

BTh2A.5 • 11:45

Optical Sensing for Temperature and Vibration Analysis of Submersible Pumps, Paulo H. Ruiz Mazzo¹, Gabriel A. Torelli¹, Uilian J. Dreyer¹, Jesse Pelegrin², Leonardo Siqueira¹, Jhoan Cubas¹, Moises Neto¹, Luiz C. Silva³, Gabriel Romero³, Jurandir Silva³, Rigoberto Morales¹, Jean Carlos Cardozo da Silva¹; ¹*Federal University of Technology-Paraná, Brazil*; ²*Instituto Federal Catarinense, Brazil*; ³*PETRÓLEO BRASILEIRO S.A. - PETROBRAS, Brazil*. Fiber Bragg Grating and Distributed Temperature Sensing are discussed to protect submersible pump components. Through a combined method using temperature and vibration measurement, we can detect early-stage issues correlated with variations in pump flow rate.

BTh2A.6 • 12:00

Small-Scale Optrode Based on PDMS for Improved Temperature Measurement, Bryan D. Sanipatin¹, Luis Sánchez², Lucía Arques¹, Salvador Sales¹; ¹*Universidad Politécnica de Valencia, Spain*; ²*Cal-Sens, Spain*. A novel, small-scale PDMS-based fiber-optic sensor is proposed to enhance sensitivity in temperature measurements. This sensor, also referred to as an optrode, was experimentally studied obtaining a substantial improvement in terms of sensitivity and size.

BTh2A.7 • 12:15

FBG-Based Accelerometer With Temperature Compensation for Structural Health Monitoring, Chloé Landreau^{2,1}, Thomas Le Gall², Nicolas Ponthus², Adriana Morana¹, Jacques Charvin², Sylvain Girard¹, Emmanuel Marin¹; ¹*Laboratoire Hubert Curien, France*; ²*Avnir Energy, France*. A Fiber Bragg Grating based accelerometer has been designed and manufactured for Structural Health Monitoring applications. It has a linear response up to 250 Hz and a temperature compensation system.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

10:30 -- 12:30

Room: 205A

ITh2B • Novel Integrated Materials II

Presider: Nathalie Vermeulen; Vrije Universiteit Brussel, Belgium

ITh2B.1 • 10:30 Invited

Wafe-Scale TMD Monolayer Waveguide, Myungjae Lee¹; ¹*Seoul National University, Korea (the Republic of)*. Abstract not available.

ITh2B.2 • 11:00

Demonstration of Graphene Waveguide Photodetector Based on Photothermoelectric Effect, Yishu Huang^{1,2}, Tom Reep^{1,2}, Hung-Chieh Tsai¹, Jeroen De Coster¹, Steven Brems¹, Inge Asselberghs¹, Klaas-Jan Tielrooij^{3,4}, Dries Van Thourhout^{1,2}, Christian Haffner¹; ¹*imec, Belgium*; ²*Ghent University, Belgium*; ³*Catalan Institute of Nanoscience and Nanotechnology (ICN2), Spain*; ⁴*TU Eindhoven, Netherlands*. We demonstrate a graphene waveguide photodetector featuring a responsivity of ~1.2 V/W. The photovoltage map measured under zero bias confirms that photothermoelectric effect is dominating.

ITh2B.3 • 11:15

Gallium Phosphide Platforms for Integrated Photonics, Lise Morice^{1,2}, Brieg Le Corre^{3,1}, Antoine Lemoine¹, Abdelmounaim Harouri³, Grégoire Beaudoin³, Luc Le Gratiet³, Tony Rohel¹, Julie Le Pouliquen¹, Rozenn Bernard¹, Christian Grillet², Charles Cornet¹, Isabelle Sagnes³, Konstantinos Pantzas³, Christelle Monat², Yoan Léger¹; ¹*Univ Rennes, INSA de Rennes, CNRS, Institut FOTON, UMR 6082, France*; ²*Institut des Nanotechnologies de Lyon, UMR CNRS 5270, Ecole Centrale de Lyon, France*; ³*Centre de Nanosciences et de Nanotechnologies, CNRS, Univ Paris-Saclay, France*. Here we compare different Gallium Phosphide photonic platforms in the framework of non-linear photonic integration. This comparison is firstly carried out through the evaluation of propagation losses within nano-waveguides in the near infrared.

ITh2B.4 • 11:30

Linear and Nonlinear Characterization of Vertical Orientation-Patterned Gallium Phosphide Waveguides for Second Harmonic Generation, Antoine Lemoine¹, Brieg Le Corre^{2,1}, Lise Morice¹, Abdelmounaim Harouri², Luc Le Gratiet², Grégoire Beaudoin², Julie Le Pouliquen¹, Arnaud Grisard³, Sylvain Combrié³, Bruno Gérard⁴, Charles Cornet¹, Yannick Dumeige¹, Konstantinos Pantzas², Isabelle Sagnes², Yoan Léger¹; ¹*Univ Rennes, INSA Rennes, CNRS, Institut FOTON - UMR 6082, F-35000 Rennes, France, France*; ²*Centre de Nanosciences et de Nanotechnologie, CNRS, 91120 Palaiseau, France, France*; ³*Thales Research and Technology, 91767 Palaiseau, France, France*; ⁴*III-V Lab, 91767 Palaiseau, France, France*. This paper discusses the potential of vertical orientation-patterned gallium phosphide (VOP-GaP) waveguide for second harmonic generation. The design of the devices, their fabrication and the linear and nonlinear characterizations are presented.

ITh2B.5 • 11:45

C-Band Translation by Second-Harmonic Generation in an Orientation-Patterned Gallium Phosphide Waveguide, Brieg Le Corre^{1,2}, Antoine Lemoine², Abdelmounaim Harouri¹, Luc Le Gratiet¹, Bruno Gérard⁴, Sylvain Combrié³, Arnaud Grisard³, Grégoire Beaudoin¹, Isabelle Sagnes¹, Konstantinos Pantzas¹, Gilles Patriarche¹, Yoan Léger²; ¹*Centre de Nanosciences et*

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

de Nanotechnologies, CNRS, France; ²Univ Rennes, INSA Rennes, CNRS, Institut FOTON - UMR 6082, France; ³Thales Research and Technology, France; ⁴III-V Lab, France. Orientation-patterned gallium phosphide (OP-GaP) is studied for frequency combs conversion from C-band to visible. Simulations of periodic and chirped OP-GaP waveguides nonlinear response are presented. Design rules for future developments are established using this study.

ITh2B.6 • 12:00 Invited

Advanced Applications of Kerr Micro-Combs, David J. Moss¹; ¹*Swinburne University of Technology, Australia.* Abstract not available.

10:30 -- 12:30

Room: 206B

NpTh2D • Topological and Quantum Nonlinear Optics

Presider: Mikko Huttunen; Tampere University, Finland

NpTh2D.1 • 10:30 Invited

Topological Optical Frequency Combs, Sunil Mittal¹; ¹*Northeastern University, USA.* We report the generation of nested topological frequency combs using two-dimensional arrays of coupled ring resonators that host robust topological edge states with linear dispersion.

NpTh2D.2 • 11:00

Topological Features of Bright Vector Solitons in Ring Resonators With Normal Dispersion, Erwan Lucas², Lewis J. Hill¹, Gang Xu³, Gian-Luca Oppo⁴, Yiqing Xu⁵, Pascal Del'Haye¹, Bertrand Kibler², Stuart Murdoch⁵, Miro Erkintalo⁵, Stephane Coen⁵, Julien Fatome²; ¹*Max-Planck-Inst Physik des Lichts, Germany; ²Laboratoire ICB, CNRS-Université de Bourgogne, France; ³School of Optical and Electronic Information, Huazhong University of Science and Technology, China; ⁴SUPA and Department of Physics, University of Strathclyde, United Kingdom; ⁵Physics Department, The University of Auckland, New Zealand.* Experiments and simulations of unusual bright vector solitons in Kerr ring resonators with normal dispersion displaying a phase defect with a phase jump of π in one of the linear polarization components are presented.

NpTh2D.3 • 11:15

Promoting Light Localization of Topological Edge States via Weak Nonlinearity, Xiaoqin Huang¹, Zhaoyuan Wang¹, Yi Hu¹, Jingjun Xu¹; ¹*Nankai University, China.* We demonstrate that a weak optical nonlinearity can further enhance the localization of topological edge states. As opposite to the reported nonlinearly-induced spreading of these states, our work introduces a solution coordinating nonlinearity and topology.

NpTh2D.4 • 11:30

Separating Spontaneous Symmetry Breaking From Exceptional Points, Lewis J. Hill¹, Julius Gohsrich^{1,2}, Jacob Fauman^{1,2}, Alekhya Ghosh^{1,2}, Kyle Kawagoe^{3,4}, Pascal Del'Haye^{1,2}, Flore Kunst^{1,2}; ¹*Max-Planck-Inst Physik des Lichts, Germany; ²Department of Physics, Friedrich Alexander University Erlangen-Nuremberg, Germany; ³Department of Physics, The Ohio State University, USA; ⁴Department of Mathematics, The Ohio State University, USA.* Spontaneous symmetry breaking is often thought of as ubiquitous with an exceptional point. However, we show that for equations describing three different systems from nonlinear optics this is probably not the case.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

NpTh2D.5 • 11:45

Non-Hermitian Swallowtail Degeneracy in the two-Mode Squeezing of Light, Polina Blinova^{1,2}, Evgeny Moiseev¹, Kai Wang¹; ¹*Physics, McGill, Canada*; ²*School of Applied and Engineering Physics, Cornell University, USA*. We show that swallowtail catastrophe consisting of various-order non-Hermitian degeneracies naturally exists in the dynamics of two-mode quadrature squeezing systems with asymmetric losses that break pseudo-Hermiticity and propose a practical experimental setup.

10:30 -- 12:30

Room: 205C

NoTh2C • 2D and Nanophotonic Devices

President: Jonathan Hu; Baylor University, United States

NoTh2C.1 • 10:30 Invited

Photoluminescence and Non-Linear Optical Behavior in Two-Dimensional Metal Chalcophosphates, Efrain Rodriguez¹; ¹*University of Maryland at College Park, USA*. Abstract not available.

NoTh2C.2 • 11:00 Invited

Non-Volatile Optical Switch Through Sliding Ferroelectricity in 2D Semiconductors, Ziliang Ye¹; ¹*University of British Columbia, Canada*. Rhombohedral stacked transition metal dichalcogenides exhibit a strong excitonic effect and an out-of-plane electrical polarization associated with the stacking configuration. Here we utilize them to achieve a non-volatile switchable behaviour in the material's optical properties.

NoTh2C.3 • 11:30 Invited

High Index Nanophotonic Structures and Optoelectronic Devices, Leland Nordin^{1,2}; ¹*CREOL, University of Central Florida, USA*; ²*Materials Science and Engineering, University of Central Florida, USA*. High-index IV-VI semiconductors promise advancements in next-generation mid-infrared devices, with high refractive index, dislocation tolerance, low Auger recombination, and cold growth temperatures. We'll discuss our heteroepitaxial IV-VI/III-V devices and high-index epitaxial nanophotonic device architectures.

NoTh2C.4 • 12:00 Invited

Transparent Conductive Oxides for Epsilon-Near-Zero and Integrated Photonics, Alan X. Wang¹; ¹*Baylor University, USA*. Transparent conductive oxides offer unique optical properties for epsilon-near-zero photonics and photonic integrated circuits. The research progress in energy-efficient optical modulators, tunable filters, and metasurfaces will be reviewed. We also discuss perspectives toward scalable manufacturing through integration with silicon photonics.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

10:30 -- 12:30

Room: 2104

SpTh2G • Signal Processing Applications

Presider: Vincent Houtsma; Nokia Bell Labs, United States

SpTh2G.1 • 10:30 Invited

Long-Haul MIMO Transmission With a Coupled Core MCF, Shohei Beppu¹, Daiki Soma¹; ¹*KDDI Research, Japan*. We review recent progress of long-haul MIMO transmission experiments with a coupled core MCF. Issues of real-time MIMO DSP implementation such as DSP complexity and adaptive tracking are also discussed.

SpTh2G.2 • 11:00

Enhanced Digital-to-Analog Converter Model Capturing Frequency Dependent ENoB, Arman Safarnejadian¹, Leslie Rusch¹, Wei Shi¹, Ming Zeng¹; ¹*ECE Department, Center for Optics, Photonics and Lasers (COPL), Canada*. We propose a digital-to-analog converter (DAC) model that accurately captures frequency-dependent effective number of bits (ENoB). Our model overcomes previous limitations (including aliasing and distortion) and reflects the DAC true frequency response.

SpTh2G.3 • 11:15 Invited

Neuromorphic Computing for Low Power DSP, Shuangxu Li¹; ¹*Huawei Technologies, Germany*. Abstract not available.

SpTh2G.4 • 11:45

Anomaly Detection in Optical Fiber: a Change-Point Detection Perspective, Reza Mosayebi¹, Lutz Lampe¹; ¹*University of British Columbia, Canada*. We present a change-point detection algorithm for optical fibers. Utilizing SNR, our approach swiftly identifies soft anomalies, aiding early failure detection. This proactive identification can mitigate connectivity disruptions, an important step toward enhancing network reliability.

SpTh2G.5 • 12:00 Invited

Fiber Sensing Using Real-Time Coherent Transceivers, Mikael Mazur¹, Nicolas K. Fontaine¹, Roland Ryf¹, Lauren Dallachiesa¹, Haoshuo Chen¹, David Neilson¹; ¹*Nokia Bell Labs, USA*. We review progress on fiber sensing using coherent transceivers and benchmark it to regular dedicated fiber sensing methods. Applications over aerial, terrestrial and transoceanic links are covered, focusing on both network protection and environmental sensing.

10:30 -- 12:30

Room: 205B

STh2F • Spectroscopy & Dynamics

Presider: Thomas Cooper; York University, Canada

STh2F.1 • 10:30 Invited

Drain and Photo-Induced Multifunctional Ambipolar Optoelectronics Based on 2D Semiconductors, Kayoung Lee¹, Jaeha Hwang¹, Youngkyu Ko¹, Suyeon Lee¹, Jungi Song¹, Yongwook Seok¹, Hanbyeol Jang², Kenji Watanabe³, Takashi Taniguchi³; ¹*Korea Advanced Institute of Science & Technology (KAIST), Korea (the Republic of)*; ²*Gwangju Institute of Science and Technology (GIST), Korea (the Republic of)*; ³*National Institute for Materials*

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Science, Japan. Practical applications of drain-bias-induced effect in the advancement of switching electronics have remained limited. Here, I will talk strategies to achieve significant current changes by utilizing drain-induced and photo-induced carrier type switching in 2D semiconductors.

STh2F.2 • 11:00

Ultrafast Hot Carrier Relaxation Dynamics in CsPbBr₃ in Presence of Layered Material, Naresh C. Maurya¹, K V Adarsh¹; ¹*IISER Bhopal, India, India.* Our study reveals that the hot carriers' initial temperature of CsPbBr₃ dropped from 1000 to 542 K and the thermalization time 550 to 350 fs in the presence of layered material, indicating hot carrier relaxation.

STh2F.3 • 11:15

Exceptionally Low-Threshold Multiple Carrier Generation in Atomically Thin Transition Metal Dichalcogenides, Riyanka Karmakar¹, Pravrati Taank¹, K V Adarsh¹; ¹*IISER Bhopal, India, India.* We experimentally demonstrate multiple carrier generation at $1.12E_g$ (E_g is bandgap) threshold energy in monolayer MoS₂ by leveraging the carrier-carrier scattering that excites secondary electrons from electron-donating sulfur-vacancy states strategically positioned below the conduction band.

STh2F.4 • 11:30 Invited

Probing Dynamics of Chemical Bonds in Organic Chromophores by X-ray Spectroscopies, Sergei Tretiak¹; ¹*Los Alamos National Laboratory, USA.* I will overview some possible measurements that can be done with X-ray lasers suggested by computational investigations. This includes monitoring the coherence evolution in molecular photoswitches and using X-ray Circular Dichroism as local chirality probe.

STh2F.5 • 12:00

Organic Photodetector Crossbar Array With Visible to Near-Infrared Responsivity, Sangin Hahn¹, Carmela Michelle Esteban¹, Sun-Woo Jo¹, Songhyun Kim¹, Kyungmin Kim¹, Seunghyup Yoo¹; ¹*KAIST, Korea (the Republic of),* This study presents the implementation of the 8x8 crossbar photodetector array, with a vertically combined organic photodiode-blocking diode pixel structure. Optimized components and simplified fabrication processes demonstrate a straightforward strategy for realizing photodetector arrays.

STh2F.6 • 12:15

Enhancing Absorption in Cadmium Telluride Solar Cells using Ti Nanoparticles, Sudarshan K. Jain¹, Nikhil Deep Gupta², Vijay Janyani¹; ¹*ECE, Malaviya National Institute of Technology, Jaipur, India;* ²*Center for VLSI and Nano Technology, Visvesvaraya National Institute of Technology, India.* This paper presents the integration of titanium nanoparticles into the active layer of CdTe solar cells to increase light absorption. It was found that titanium nanoparticles' light-trapping capability profoundly improved the performance of the CdTe solar cells.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

10:30 -- 12:30

Room: 207

SoTh2E • Fibers and Devices for Biomedical and Sensing Applications II

Presider: Natasha Vukovic; University of Southampton, United Kingdom

SoTh2E.1 • 10:30 Invited

Bidirectional Interfaces Based on Multimaterial Optical Fibers for Infrared

Neurostimulation, Marcello Meneghetti^{1,2}, Kunyang Sui^{1,2}, Rune W. Berg², Christos Markos^{1,3};

¹*Technical University of Denmark, Denmark*; ²*University of Copenhagen, Denmark*; ³*NORBLIS Aps, Denmark*. Here, we present the development of multifunctional neural implants based on soft optical fibers, and their application to infrared neural stimulation, a transgene-free neuromodulation technique with high potential for clinical translation.

SoTh2E.2 • 11:00 Invited

Multifunctional Neural Interfaces With Multimodal Optical Fibers, Ferruccio Pisanello¹,

Marco Bianco¹; ¹*Istituto Italiano di Tecnologia, Italy*. Abstract not available.

SoTh2E.3 • 11:30

Thermally Drawn Biodegradable Optical Fiber for Neural Applications, Parinaz

Abdollahian^{1,2}, Kunyang Sui^{1,2}, Guanghui Li², Jiachen Wang¹, Cuiling Zhang¹, Yazhou Wang¹, Rune W. Berg², Marcello Meneghetti^{1,2}, Christos Markos¹; ¹*Photonics and Electronic Engineering, Technical University of Denmark, Denmark*; ²*Neuroscience, Copenhagen University, Denmark*. This study focuses on the development of implantable, biodegradable optical fibers for light delivery in the brain without the necessity of explanation surgeries. The biodegradability of the fibers was verified *in vitro* and *in vivo*.

SoTh2E.4 • 12:00

Ultra-High Transverse Mode Purity in Double-Clad Hollow-Core Photonic Crystal Fiber,

Zhuozhao Luo^{1,2}, Jiapeng Huang^{1,3}, Yu Zheng⁴, Ruochen Yin^{1,3}, Long Zhang¹, Meng Pang^{1,2}, Xin Jiang^{1,3}; ¹*Russell Centre for Advanced Lightwave Science, Hangzhou Institute of Optics and Fine Mechanics and Shanghai Institute of Optics and Fine Mechanics, China*; ²*State Key Laboratory of High Field Laser Physics and CAS Center for Excellence in Ultra-intense Laser Science, Shanghai Institute of Optics and Fine Mechanics, Chinese Academy of Sciences, China*; ³*Innovation and Integration Center of New Laser Technology, Shanghai Institute of Optics and Fine Mechanics, Chinese Academy of Sciences, China*; ⁴*Fiber Optoelectronics Technology Co., Ltd., China*. Ultra-high single-mode purity is achieved in a double-clad hollow-core photonic crystal fiber by enhanced higher-order-mode filtering. The wavelength tunability with persisted high single-mode purity is also investigated.

14:00 -- 16:00

Room: 206A

BTh3A • FBG for Laser and Spectrometer Applications

Presider: Sébastien Loranger; Polytechnique Montréal, Canada

BTh3A.1 • 14:00 Invited

Beyond 2 μm Bragg Grating Components for Monolithic Fiber Lasers, Antreas

Theodosiou¹, Jan aubrecht², Pavel Peterka², Kyriacos Kalli³, Ori Sapir-Henderson⁴; ¹*Lumoscribe Ltd., Cyprus*; ²*UFE, Czechia*; ³*CUT, Cyprus*; ⁴*UoA, Australia*. We report on our latest results on

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

the inscription of uniform and blazed fiber Bragg gratings in fluoride and silica optical fibers for monolithic SWIR fiber lasers.

BTh3A.2 • 14:30

Femtosecond Written Chirped Fiber Bragg Gratings for Dispersion Control at 2 μm ,

Georg R. Schwartz¹, Ria G. Krämer¹, Malte P. Siems¹, Abdolnaser Ghazagh¹, Daniel Richter¹, Stefan Nolte^{1,2}; ¹*Institute of applied physics, Friedrich-Schiller-Univ., Germany*; ²*Fraunhofer Institute for Applied Optics and Precision Engineering IOF, Germany*. We present femtosecond written CFBGs using a chirped phase mask for dispersion control. A bandwidth of >140 nm and reflectivity of ~70 % were achieved, with tailored dispersion up to the 3rd order.

BTh3A.3 • 14:45

High Efficiency Fiber Bragg Grating Spectrometer Fabricated Using One Infrared

Femtosecond Laser Pulse and the Phase Mask Technique, Abdullah Rahnama¹, Cyril Hnatovsky¹, Robert B. Walker¹, Kasthuri De Silva¹, Stephen J. Mihailov¹; ¹*National Research Council Canada, Canada*. A super-efficient all-fiber visible spectrometer with sub-nanometer resolution is fabricated using one infrared femtosecond laser pulse and a phase mask. The strong light outcoupling from the fiber is due to micropores formed in its core.

BTh3A.4 • 15:00

Selective Modal Excitation of FBGs in FMF Through Inscription Techniques and the use

of a Spatial Multiplexer, James T. Hainsworth^{1,2}, Adriana Morana², Lucien Pouget¹, Marina Arnaud¹, Sylvain Girard², Jacques Decroix¹, Emmanuel Marin²; ¹*IRT Saint Exupery, France*; ²*Laboratory Hubert Curien, France*. Fibre Bragg grating spectral control by use of a spatial mode sorter and inscription eccentricity has been investigated. Results show extensive peak suppression in the transmission and reflection spectra with the capability for peak erasure.

BTh3A.5 • 15:15

Chirped Tilted Fiber Bragg Gratings Used as Intracavity ASE Band-Stop Filters in a High

Power 1018 nm Fiber Laser, Bertrand Morasse¹, Alexandre Perron¹, Dominic Faucher¹, Pierre-Michel Belzile¹, Mathieu Vollant Deschênes¹, Frédéric Faucher¹, Guillaume Brochu¹, François Trépanier¹, Michel Bégin¹, Pascal Deladurantaye¹; ¹*TeraXion Inc., Canada*. We present chirped tilted fiber Bragg grating intracavity filters to suppress amplified spontaneous emission from the core of a 20/400 Yb-doped high power fiber laser, allowing the efficient generation of 432 W at 1018 nm.

BTh3A.6 • 15:30

Effect of Group Delay Ripples of Chirped Fiber Bragg Gratings Used in CPA Lasers,

Francois Ouellette¹, Hui Wang²; ¹*Chengdu University, China*; ²*Ultron Photonics, China*. The effect of group delay ripples (GDR) in chirped fiber Bragg gratings used in chirped pulse amplification lasers is analyzed. The polarization rotation method is shown to measure the most deleterious GDR with a resolution better than 50 fs.

BTh3A.7 • 15:45

Femtosecond-Written Large Area FBGs for Wavelength Stabilization of Blue Laser

Diodes, Ludovic de Repentigny^{2,1}, Lauris Talbot², François Trépanier¹, Martin Bernier²; ¹*TeraXion Inc., Canada*; ²*Centre d'Optique, Photonique et Laser (COPL), Université Laval,*

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Canada. We report the wavelength stabilization of blue laser diodes using fiber Bragg gratings femtosecond-written in their highly multimode fiber pigtail.

14:00 -- 16:00

Room: 206B

NpTh3C • Novel Nonlinear Materials

Presider: Mikko Huttunen; Tampere University, Finland

NpTh3C.1 • 14:00

An Ultrafast All-Optical Switch With an Epsilon-Near-Zero-Based Nanocavity, Yaswant Vaddi¹, Theng-Loo Lim¹, M. Zahirul Alam¹, Shivashankar R. Vangala², Jeremy Upham¹, Joshua Hendrickson², Robert W. Boyd^{1,3}; ¹*Department of Physics, University of Ottawa, Canada;* ²*Sensors Directorate, Air Force Research Laboratory, USA;* ³*Institute of Optics, University of Rochester, USA.* We experimentally demonstrate an ultrafast all-optical switch using a 1D, non-linear nanocavity with an epsilon-near-zero mirror. The switch exhibits a 10 dB modulation depth over a large spectral range.

NpTh3C.2 • 14:15

Tailoring Nonlinear Response of ENZ Metamaterials at Oblique Incidence, Sisira S. Suresh¹, M. Zahirul Alam¹, Jeremy Upham¹, R. Margoth Cordova Castro¹, Maryam Abbasi¹, Robert W. Boyd¹; ¹*University of Ottawa, Canada.* We study nonlinear optical properties of a layered ENZ metamaterial with oblique incident TM-polarized light. Maximum nonlinear responses occur at angle-dependent zero-permittivity wavelengths, demonstrating enhancement is due to ENZ condition regardless of constituents' microscopic details.

NpTh3C.3 • 14:30

Enhanced THz Third-Harmonic Generation in a Graphene-Metamaterial Hybrid Structure, Ali Maleki¹, Moritz B. Heindl², Yongbao Xin³, Robert W. Boyd^{1,4}, Georg Herink², Jean-Michel Ménard¹; ¹*University of Ottawa, Canada;* ²*University of Bayreuth, Germany;* ³*Iridian Spectral Technologies Ltd, Canada;* ⁴*University of Rochester, USA.* We present a metasurface design to enhance third harmonic generation at terahertz frequencies inside two decoupled graphene sheets. Spectral filtering of terahertz pulses produced by a table-top source enables sensitive monitoring of nonlinear effects.

NpTh3C.4 • 14:45

All-Dielectric Huygens' Metawaveguides for Nonlinear Integrated Photonics, Gabriel Flizikowski¹, Ozan W. Oner², Lais F. dos Santos¹, M. Saad Bin-Alam³, Thomas Pertsch⁴, Isabelle Staude⁴, Jens H. Schmid³, Pavel Cheben³, Ksenia Dolgaleva^{1,2}; ¹*School of Electrical Engineering and Computer Science, University of Ottawa, Canada;* ²*Department of Physics, University of Ottawa, Canada;* ³*National Research Council of Canada, Canada;* ⁴*Friedrich-Schiller-Universität Jena, Germany.* We have experimentally demonstrated the compensation of nonlinear optical effects by a silicon-on-insulator nanophotonic metawaveguide comprising a chain of resonantly forward-scattering nanoparticles, exhibiting soliton-like behavior and spectral pulse integrity.

NpTh3C.5 • 15:00

Highly Efficient Degenerate Four-Wave Mixing With an Epsilon-Near-Zero-Based Low-Q Cavity, Theng Loo Lim¹, Yaswant V. Vaddi¹, M. Zahirul Alam¹, Shivashankar R. Vangala³,

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

Jeremy Upham¹, Joshua Hendrickson³, Robert W. Boyd^{1,2}; ¹*University of Ottawa, Canada*; ²*University of Rochester, USA*; ³*Air Force Research Laboratory, USA*. We experimentally demonstrated high-efficiency frequency-degenerate four-wave mixing (DFWM) using an epsilon-near-zero-based nanocavity. The cavity length is approximately two wavelengths thick. We measured an absolute efficiency as large as 34%.

NpTh3C.6 • 15:15

Second Harmonic Enhancement With a U-Shaped InSb Plasmonic Antenna in the THz Regime, Sina Aghili¹, Ksenia Dolgaleva¹; ¹*University of Ottawa, Canada*. We propose a U-shaped plasmonic antenna made of the InSb semiconductor, enhancing the SHG process with an effective second-order susceptibility of 90 pm/V in the THz regime

NpTh3C.7 • 15:30 Invited

Nonlinear Laser Conversion in Nanophotonic Resonators, Jennifer A. Black¹; ¹*NIST Boulder, USA*. Integrated Kerr microresonators incorporating nanostructures provide a novel approach to phase-matching for flexible and efficient laser wavelength conversion based on optical-parametric oscillation. I will discuss the development of these nanophotonic oscillators for laser wavelength conversion spanning beyond an octave.

14:00 -- 16:00

Room: 205C

NoTh3B • Emerging Photonic Devices

Presider: Lynda Busse; US Naval Research Laboratory, United States

NoTh3B.1 • 14:00 Invited

Near-Ultraviolet to Midwave Infrared Devices for Quantum Sensing and Information Processing, Cheryl M. Sorace-Agaskar¹, Colin Bruzewicz¹, Patrick Callahan¹, Christopher Heidelberger¹, Dave Kharas¹, William Loh¹, Thomas Mahony¹, Ryan Maxson¹, Robert McConnell¹, Alexander Medeiros¹, Rachel Morgan², Alkesh Sumant¹, Meghan Schuldt¹, Reuel Swint¹, Kerri Cahoy², Jelena Notaros², John Chiaverini^{1,2}, Paul Juodawlkis¹; ¹*MIT Lincoln Laboratory, USA*; ²*MIT, USA*. The talk reviews photonic integrated circuit materials, devices and integration techniques developed at MIT Lincoln Laboratory to support the needs of next generation quantum systems across the wavelength spectrum from the near-ultraviolet to the midwave-infrared.

NoTh3B.2 • 14:30 Invited

Deep Ultraviolet MicroLEDs and Arrays for Advanced Optical Communication, Haiding Sun¹; ¹*University of Sci. & Tech. of China, China*. Abstract not available.

NoTh3B.3 • 15:00 Invited

Harnessing Ultrafast Optical Pulses for 3D Microfabrication by Selective Tweezing and Immobilization of Colloids, Krishangi Krishna¹, Jieliyue Sun¹, Kimani C. Toussaint¹; ¹*Brown University, USA*. We employ femtosecond-laser pulses for a two-step microfabrication process: optical tweezing of microparticles to target positions, followed by immediate particle immobilization by two-photon polymerization. The challenges and opportunities of this platform are discussed.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

NoTh3B.4 • 15:30

Withdrawn

NoTh3B.5 • 15:45

Mie-Resonant Huygens' Metawaveguide Microring Resonators, Md Saad-Bin-Alam¹, Jianhao Zhang¹, Yunus Denizhan Sirmaci², Thomas Pertsch², Isabelle Staude², Jens H. Schmid¹, Pavel Cheben¹; ¹*National Research Council Canada, Canada*; ²*Friedrich-Schiller-Universität Jena, Germany*. Metawaveguides consisting of Mie-resonant dielectric Huygens' nanoantennas can propagate light with negative group index and zero-to-anomalous dispersion. Here, we demonstrate a Huygens' metawaveguide microring resonator, which could revolutionize the on-chip nonlinear and quantum photonics applications.

14:00 -- 16:00

Room: 207

SoTh3D • Novel Fabrication & Characterization Techniques

Presider: Nadia Boetti; *Fondazione LINKS, Italy*

SoTh3D.1 • 14:00 Invited

X-ray-Based Techniques for the Characterization of Polymer Optical Fibers, Mario Ferraro^{1,3}, Maria Caterina Crocco^{1,3}, Raffaele Filosa^{1,3}, Kevin Kiedrowski², Marco Jupé², Marco Leonetti⁴, Giancarlo Ruocco⁴, Stefan Wabnitz⁵, Riccardo C. Barberi^{1,3}, Vincenzo Formoso^{1,3}, Raffaele G. Agostino^{1,3}; ¹*STAR Research Infrastructure, University of Calabria, Italy*; ²*Laser Zentrum Hannover e.V., Germany*; ³*Physics Department, University of Calabria, Italy*; ⁴*Center for Life Nano Science@Sapienza, Italian Institute of Technology, Italy*; ⁵*DIET, University of Rome "La Sapienza", Italy*. We report on a characterization study of polymer optical fibers based on X-ray computed microtomography and X-ray small-angle scattering.

SoTh3D.2 • 14:30 Invited

Silica Specialty Fibers Made Through Laser-Assisted Additive Manufacturing, Pawel Maniewski^{1,2}, Valdas Pasiskevicius¹, Christopher Holmes²; ¹*KTH Royal Institute of Technology, Sweden*; ²*Optoelectronic Research Centre, University Of Southampton, United Kingdom*. Novel approaches for laser-based silica processing are demonstrated, that offer unique fabrication capabilities for specialty fibers. High performance and new fiber geometries are offered through multi-material additive manufacturing, cutting, polishing, welding, and laser-based preform drawing.

SoTh3D.3 • 15:00

Systematic Study of Flame-Fabricated Surface Nanoscale Axial Photonics Resonators, Asaad Hannah¹, Emily Eadie¹, Samar Deep¹, Pablo Bianucci¹; ¹*Concordia University, Canada*. We present a systematic investigation of the different parameters involved in the fabrication of surface nanoscale axial photonics resonators from optical fibers using a flame, and how they affect the resulting resonant mode structure.

SoTh3D.4 • 15:15

Accurate Measurement of Wavelength-Dependent Beam Parameters of a Supercontinuum Laser Source Focused by a Lensed Fiber Probe, Kuan-Yuan Chang¹, Jun-Cheng Hsu¹, Jia-Ming Liu^{1,2}; ¹*National Yang Ming Chiao Tung University, Taiwan*; ²*Electrical and Computer Engineering, University of California, Los Angeles, USA*. A supercontinuum (SC)

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

laser is focused by an optical lensed fiber (OLF). By applying a spectrometer to measure back-coupling efficiency, multi-wavelength parameters of the OLF probe and the focused SC laser beam can be analyzed.

SoTh3D.5 • 15:30

Enhanced Ytterbium Emission – Manifold Resolution at Room Temperature Using the Purcell Effect in Doped Silica Microspheres, Nikita Toropov¹, Christophe A. Codemard², Neil P. Sessions¹, Michalis N. Zervas¹; ¹*University of Southampton, United Kingdom*; ²*TRUMPF Lasers UK Ltd, United Kingdom*. We propose and demonstrate a novel method to characterize the Stark-split emission manifold of rare-earth doped silica glass at room temperature, using the Purcell effect in Yb-doped microspheres.

16:30 -- 18:30

Room: 2000A

JTh4A • Joint Postdeadline Paper Presentations and Student Award Winner Announcement

JTh4A.1 • 16:30 Postdeadline Submission

High Fiber-to-Fiber Gain low Noise Figure Erbium Doped Waveguide Amplifiers in the Manufacturable Al₂O₃ Platform, Carlos E. Osornio-Martinez¹, Dawson B. Bonneville¹, Meindert Dijkstra¹, Sonia Garcia-Blanco¹; ¹*Universiteit Twente, Netherlands*. Erbium doped waveguide amplifiers (EDWAs) with performance approaching fiber amplifiers (i.e., net fiber-to-fiber gain (~17 dB), noise figure (~6 dB) and off-chip output power (~15 dBm) in the manufacturable Al₂O₃ integrated photonics platform are reported.

JTh4A.2 • 16:45 Postdeadline Submission

Microstructured Optical Fibers Made by Additive Manufacturing of Chalcogenide Glasses, Johann Troles¹, Leo Szymczyk^{1,2}, Francois Chevire¹, Catherine Boussard¹, Antoine Gautier¹, Frederic Charpentier², Yann Guimond², Mathieu Roze², Gilles Renversez³; ¹*University of Rennes, France*; ²*UMICORE IR Glass, France*; ³*Centrale Marseille, France*. Chalcogenide preforms obtained by an additive manufacturing process have been drawn into chalcogenide optical fibers. Those results open a way for the elaboration of chalcogenide microstructured optical fibers, especially for hollow-core fibers.

JTh4A.3 • 17:00 Postdeadline Submission

Four-Fold Truncated Double-Nested Anti-Resonant Hollow-Core Fiber for Ultralow Loss and Robust Single Mode Operation, Shoufei Gao^{2,1}, Yizhi Sun^{1,2}, Hao Chen¹, Dawei Ge³, Dong Wang³, Dechao Zhang³, Han Li³, Wei Ding^{1,2}, Yingying Wang^{2,1}; ¹*Jinan University, China*; ²*Linfiber Technology (Nantong) Co., Ltd., China*; ³*China Mobile Research Institute, China*. We report the simultaneous achievement of ultralow loss in fundamental mode (0.1-0.2 dB/km) and high loss in higher order modes (db/m level) with an extinction ratio reaching 26000 in a novel four-fold truncated DNANF structure.

JTh4A.4 • 17:15 Postdeadline Submission

Degenerate Optical Parametric Oscillation in Coupled Cavities, Laís Fujii dos Santos¹, Felipe G. Santos², Ksenia Dolgaleva¹; ¹*University of Ottawa, Canada*; ²*Quantum Valley Ideas Laboratories, Canada*. Networks of coupled parametric oscillators exhibiting binary phase state

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

can be used for computation. The phase relation between oscillators can be switched with frequency detuning.

JTh4A.5 • 17:30 Postdeadline Submission

Machine Learning-Assisted Extreme Events Forecasting, Saliya Coulibaly¹; ¹*Universite de Lille, France*. Here, we propose a new supervised machine learning strategy to locally forecast bursts occurring in the turbulent regime of a fiber ring cavity.

JTh4A.6 • 17:45 Postdeadline Submission

Radiation Hardness Evaluation of Anti-Resonant Hollow Core Fibers for Extreme Environments, Amy Van Newkirk¹, William Lo², Matt Leoschke², Marcello Catellani², Michael Reilly¹, J. Enrique Antonio Lopez³, Rodrigo Amezcua Correa³, Axel Schulzgen³, Shlomi Zilberman², Federico Scurti²; ¹*Penn State Electro-Optics Center, USA*; ²*Ken and Mary Alice Lindquist Department of Nuclear Engineering, Pennsylvania State University, USA*; ³*CREOL, University of Central Florida, USA*. An ARHCF was irradiated with neutron and gamma radiation from a nuclear reactor. The RIA was 5x less than that of pure silica core fiber, showing significant promise for applications in extreme radiation environments.

Optica Advanced Photonics Session Guide

Disclaimer: this guide is limited to technical program with abstracts and author blocks as of 21 July. For updated and complete information with special events, reference the online schedule or mobile app.

JD1 • Joint On-demand Session

JD1.1

Low-Cost Egg White Based Whispering Gallery Mode Micro-Laser, Yang Luo¹, Jun-Hua Huang¹, Jian-Kun Wang¹, Geng-Liang Chen¹, Guo-Liang Zheng¹, Xuhui Zhang¹; ¹*Shenzhen Technology University, China*. We present a protein microsphere laser by dehydration by mixing egg white with dye solution, and analyze the data of its optical characterization measurement, which verifies that it has great potential for application in bio-microlaser.

JD1.2

Design of Colored Fluorescent Radiative Cooling Bilayer Polymer Coatings by Modified Monte Carlo Method, Tao Wang¹, Danyuan Lei², Jianguo Dai²; ¹*The Hong Kong Polytechnic University, Hong Kong*; ²*City University of Hong Kong, Hong Kong*. Colored radiative cooling is challenging. Here, a modified Monte Carlo method was proposed for design of colored fluorescent radiative cooling coatings. Colored bilayer coatings were designed for efficient cooling performance.

JD1.3

Enhanced Nonlinearity in VSe₂-SWCNT Nanohybrid, Vinod Kumar¹, Chandra S. Rout², K. V. Adarsh¹; ¹*IISER Bhopal, India*; ²*Jain university, India*. We demonstrate enhanced third-order nonlinear optical response in the charged-coupled VSe₂-SWCNT hybrid attributed to charge transfer. This enhancement paves the way for potential applications in optical limiting, with an impressive onset threshold of 40 GW/cm².